



BIOECOLOGY AND MANAGEMENT OF MAJOR INSECT PESTS OF RICE IN WEST BENGAL

S DUTTA¹, A HOSSAIN^{2,3} AND N ROY^{4*}

¹Rabindra Mahavidyalaya, Department of Zoology, Champadanga, Hooghly 712401, West Bengal, India

²Ecological Research Lab, Sidho Kanho Birsa University, Department of Zoology, Purulia 703104, West Bengal, India

³Department of Zoology, The University of Burdwan, Golapbag, Burdwan 713104, West Bengal, India

⁴Ecology Research Unit, MUC Women's College, Department of Zoology, Burdwan 713104, West Bengal, India

*Email: nayan909@gmail.com (corresponding author): ORCID ID 0000-0002-0114-0596

ABSTRACT

West Bengal is considered a large producer state of rice in India with an output of nearly 55.48% of total crop production of Bengal. In West Bengal, rice has grown in three different cropping seasons such as Aus (autumn rice), Aman (winter rice) and Boro (summer rice), contributing about 20.69% to the total net State Domestic Product (SDP). Rice production in West Bengal hampered by many biotic stresses mostly by stem borers [*Scirphophaga incertulas* (Walker, 1863), *Sesamia inferene* (Walker, 1856), *Chilo polychrysus* (Meyrick, 1932), *Chilo suppressalis* (Meyrick, 1863)] leaf folder [*Cnaphalocrocis medinalis* (Guenee, 1854)], plant hoppers [*Nilaparvata lugens* (Stal, 1854), *Sogatella furcifera* (Horvath, 1899)], leaf hoppers [*Nephotettix virescens* (Distant, 1908), *Nephotettix nigropictus* (Stal, 1870)], rice hispa [*Diclidispa armigera* (Olivier, 1808)], rice bugs [*Leptocorisa acuta* (Thunberg, 1783), *Leptocorisa oratorius* (Fabricius, 1794), *Brevennis rehi* (Lindinger, 1943)], gall midge [*Orseolia oryzae* (Wood-Mason, 1889)], etc. Demographic parameter has shown that rice pests are greatly affecting rice host either caused directly by creating dead heart (DH), hopperburn, onion leaf, silver shoot during different stages of harvesting or by indirectly transmitting grassy stunt virus (RGSV), rice ragged stunt virus (RRSV), rice tungro viruses (RTV), rice tungro bacilliform virus (RTBV) etc. resulting 9.5% production loss. For minimize the infestation farmers use different rice varieties which provide resistance effect in pest population. For controlling rice pest population farmers mostly use chemical insecticide [e.g. fipronil, acephate, fipronil, monocrotophos] which exhibit good results against rice pests. Beside those other techniques including using biological and biochemical pesticides [neem oil, karanj oil, mahua oil, conidial suspension of *M. anisopliae*], pheromones [(Z)-11-hexadecinal, (Z)-9-hexadecenal], allomones [5: 1 mixture of 2-(E)-octenyl acetate and octanol 3-octenal], cultural methods [synchronized planting, using of trap crop etc.] are effective for controlling rice pests. The population ecology based sustainable management of such pest species will support E³ strategy [Ecosystem service-based ecological engineering for ecological pest management (ESS-EE-EPM)] of pest management for successful cultivation of rice in the near future.

Key words: *Oryza sativa*, state domestic product (SDP), West Bengal biotic stresses, sustainable management, E³ strategy, chemical, pests, status, cultural, biopesticides, biological resistant variety, pheromones, population based control

Rice (*Oryza sativa* L., Gramineae or Poaceae, 2n=24) continues to remain as the staple food of West Bengal by contributing approximately 55.48% of total crop production (Adhikari et al., 2012; Bag et al., 2011; Dey et al., 2005). Several limiting factors such as flood, pest infestation, damage by animal and human activity causing yield loss thereby narrow down the profit of Bengal farmer (Dutta and Roy, 2022; Bag et al., 2011). Among them insect pests have appeared as the most important limiting factors during pre- and post-harvest stage of rice cultivation (Arora et al., 2019;

Adhikari et al., 2019). For controlling rice insect pest in Bengal, farmers have been most frequently using synthetic pesticides in broad-spectrum which shown good efficacy against herbivorous insect pests (Adhikari et al., 2019; Tigga et al., 2018). Besides that, using of biochemical pesticides, biological agents, pheromone trap, repellent allomone, cultural strategies and several other modern strategies like selection of high yielding resistant varieties, sterile male technique by irradiation or genetically engineered pests in management of respective pest species in the field also shown good result

for controlling rice insect pests (Sankar and Rani, 2018; Gunawardena and Bandumathie, 1993; Padhan and Raghuraman, 2018). All of these management processes have imparted negative impact also (Chatterjee et al., 2008). Sometime pest population create resistance against applied pesticide and also these pesticide targets beneficial faunal biodiversity (pollinators, natural enemy of pest etc.) which create destabilizing agro ecosystem (Keary, 2023; Roy, 2020, 2021, 2022). Beside it has observed that all of rice pest management techniques did not consider economic threshold levels (ETLs) limit for pest that create ecological imbalance (Carvalho, 2017; Mobarak et al., 2020). These result into secondary pest outbreak, pest resurgence, and emergence of new pest biotypes, destabilized prey (rice pest)-predator (natural enemy of rice pest) stability, which ultimately leads regulatory complications with several constraints and limitations on the agro ecosystem (Dutta and Roy, 2016; Kakde and Patel, 2018; Kakde et al., 2014; Jeevanandham et al., 2020; Chávez et al., 2018; Roy, 2020, 2021). Therefore, this study emphasizes that the life cycle of major rice pests, their active and passive infestation and their management that will be very helpful to point out a possible scope for sustainable ecological management towards climate smart pest management (CSPM) through supporting E³ strategy [Ecosystem service-based Ecological engineering for Ecological pest management (ESS-EE-EPM)] of pest management for successful cultivation of rice crop in the near future.

Status of cultivation

Agriculture is the leading occupation of West Bengal and rice is the state's principal food crop (Chatterjee et al., 2008; Adhikari et al., 2012). Rice cultivation in West Bengal has occurred in diversified ecosystems such as irrigated, upland, rainfed low land, semi deep water and deep water as well as coastal saline regions in three different cropping seasons, [Aus (autumn rice), Aman (winter rice) and Boro (summer rice)] (Dhaliwal et al., 2010; Azgar and Hembram, 2018; Aryal et al., 2018). Rice was cultivated 61% of the total geographical area in West Bengal, contributing about 20.69% of State Domestic Product (Adak et al., 2020). Rice in West Bengal generally cultivated in three different seasons Boro, Aus and Aman crop season (Dutta and Roy, 2022; Bag et al., 2011). Six different agro-climatic zones (Northern Hill Zone, Terai-Teesta Alluvial Zone, Gangetic Alluvial Zone, Red and Laterite Zone, Vindhya Alluvial Zone, Coastal Saline Zone) of West Bengal is a producer of high yielding (Annada, Jamini, Bhupen, IR 36, IR 64, and IR 22 etc.), traditional or local

(Bhadoo, Kabiraj Sal, and Shatia etc.), and aromatic rice varieties [Basmati, Badsabhog, Gobindabhog, Tulaipanji, etc.] (Bag et al., 2011; Mandal et al., 2022; Chatterjee et al., 2008; Dutta and Roy, 2022). This rice production was mostly hampered by insect pest infestation, accounted 7.37% of total rice production (Ghosh et al., 2015; Aryal et al., 2018) (Table 1, 2).

Status of rice insect pests

At the present agricultural scenario, loss in production of rice crop is due to a variety of major insect pest infestation (Chakraborty and Deb, 2012; Ghule et al., 2008). Generally, the crop loss at the farm level during pre and post harvest level occurred by major insect pests like stem borers, gall midge, plant hoppers, leaf folders, rice hispa, and rice bug etc. (Sarkar et al., 2016; Chakraborty and Deb, 2012) (Table 1, 2).

A. Stem borers

The rice production in West Bengal mostly hampered by infestation causing Lepidopteran stem borer species which include yellow stem borer (YSB), *Scirphophaga incertulas* (Walker, 1863) (Lepidoptera: Pyralidae), pink stem borer (PSB), *Sesamia inferens* (Walker, 1856) (Lepidoptera: Pyralidae), striped stem borer (SSB) *Chilo suppressalis* (Meyrick, 1863) (Lepidoptera: Pyralidae), dark headed stem borer (DHSB) *Chilo polychrysus* (Meyrick, 1932) (Lepidoptera: Pyralidae), (Shivasharanappa et al., 2010; Dutta and Roy, 2022; Chakraborty, 2012). *Scirphophaga incertulas* (Walker) encompasses egg, 5-6 instars larvae, pre-pupa, pupa and adult during their life cycle (Hamsein et al., 2020; Dutta and Roy, 2018b). *Sesamia inferens* (Walker) encompasses egg, 6-8 instars larvae, pre-pupa, pupa and adult during their life cycle (Baladhiya et al., 2018; Jeevanandham et al., 2020). *Chilo suppressalis* (Meyrick) has passed egg, 4 instars larvae, pupa to become full grown adult (Pathak and Khan, 1994; Dutta and Roy, 2022). *Chilo polychrysus* (Meyrick) also shown same development pattern except 3 instars larva has shown instead of 4 instars of larva (Bhatt et al., 2018; Luo et al., 2016). Damage caused by stem borer has encountered due to internodal penetration by newly hatched larva during vegetative phase specially in booting or flowering stage which remain inside or behind leaf sheath in groups and feed on the epidermal layer of leaf sheath formation of oblong holes in parallel rows in the unfolded leaves as a result central leaf turns brownish, a condition known as 'Dead Heart' (DH) (Bhatt et al., 2018; Pathak and Khan, 1994). When infestation was severe resulting folding of older rice

Table 1. Biological parameters and host range of different rice pests

Rice pest	Common name and Scientific name	Stages in life cycle	Parameters of life cycle	Period	Reference
Stem borer	Yellow stem borer <i>Scirphophaga incertulas</i>	Egg, larvae (5-6 instars), pre-pupa, pupa and adult	Oviposition period	26.0 hours	Hamsein et al., 2020; Dutta and Roy, 2018b; 2022.
			Larval period	20-40 days	
	Pink stem borer <i>Sesamia inferenze</i>	Egg, larvae (6-8 instars), pre-pupa, pupa and adult	Pre-pupal period	1-2 days	Baladhiya et al., 2018; Jeevanandham et al., 2020; Dutta and Roy, 2022.
			Pupal period	6-12 days	
			Number of days in life cycle	52-64 days	
			Number of generation each year	5-8	
			Oviposition period	2 days	
			Larval period	21-39 days	
	Stripped stem borer <i>Chilo suppressalis</i>	Egg, larvae (4 instars), pupa and adult	Pre-Pupal period	1-2 days	Pathak and Khan, 1994; Dutta and Roy, 2022; Bhatt et al., 2018.
			Pupal period	8-14 days	
Number of days in life cycle			45-57 days		
Number of generation each year			4-5		
Leaf folder <i>Cnaphalocrocis medinalis</i>	Oviposition period	Egg, larvae (3 instars), pupa and adult	Number of days in life cycle	5 days	Dutta and Roy, 2022; Bhatt et al., 2018; Luo et al., 2016.
			Number of generation each year	6	
	Larval period	Egg, larvae (5-6 instars), pupa, and adult	Oviposition period	3 days	Dutta and Roy, 2022; Bhatt et al., 2018; Luo et al., 2016.
			Larval period	23-39 days	
	Pupal period	Egg, larvae (3 instars), pupa and adult	Pupal period	7 days	Dutta and Roy, 2022; Bhatt et al., 2018; Luo et al., 2016.
			Number of days in life cycle	35-60 days	
	Leaf folder <i>Cnaphalocrocis medinalis</i>	Egg, larvae (5-6 instars), pupa, and adult	Number of generation each year	2	Bhatt et al., 2018; Gangwar, 2015.
			Oviposition period	3 days	
			Larval period	14-23 days	
			Pupal period	6-10 days	
Number of days in life cycle			15-28 days		
Number of generation each year			11		
Brown planthopper <i>Nilaparvata lugens</i>	Egg, Five nymphal stage and Adult	Oviposition period	10 days	Catindig et al., 2009; Bottrell and Schoenly, 2012.	
		Nymphal period	18-32 days		
		Number of days in life cycle	25 to 44 days		
		Number of generation each year	12-13		
		Oviposition period	2 days		
White backed planthopper <i>Sogatella furcifera</i>	Egg, five nymphal instar, adult	Nymphal period	15-28 days	Kaur et al., 2015; Kumar, 2019.	
		Number of days in life cycle	20-31 days		
		Number of generation each year	5-6		

(contd.)

(Table 1 contd.)

Leafhopper	Green leafhopper <i>Nephotettix virescens</i> and <i>Nephotettix nigropictus</i>	Egg, five instars nymph and adult	Oviposition period Nymphal period Number of days in life cycle Number of generation each year	9-14 days 12-18 days 21-35 days 11	Bunawan et al., 2014; Bhatt et al., 2018; Dey, 2016.
	Rice hispa <i>Dicladispa armigera</i>	Egg, four instars larva, pupa, adult	Oviposition period Larval period Pupal period Number of days in life cycle Number of generation each year	12 days 11-14 days 7-8 days 18-36 days 6	Iqbal et al., 2015, Bhattacharjee and Ray, 2017.
Rice bug	Gundhi bug <i>Leptocorisa oratorius</i>	Egg, five nymphal stage and adult	Oviposition period Nymphal period Number of days in life cycle Number of generation each year	12 days 24-36 days 24-36 days 2-5	Khare and Prakash, 2020; Arora et al., 2019.
	Earhead bug <i>Leptocorisa acuta</i>	Egg, Six nymphal stage and Adult	Oviposition period Nymphal period Number of days in life cycle Number of generation each year	5 days 22-24 days 30 to 60 days 4-5	Dutta and Roy, 2016, 2018a; Pathak and Khan, 1994; Bhatt et al., 2018.
	Rice mealy bug <i>Brevinnia rehi</i>	Egg, Four nymphal instars and Adult	Oviposition period Nymphal period Number of days in life cycle Number of generation each year	26 hours 14-30 days 17-37 days 12-13	Bhatt et al., 2018; Pathak and Khan, 1994.
Gall midge	Rice gall midge <i>Orseolia oryzae</i>	Egg (3 instar), larva, pupa and adult	Oviposition period Larval period Pupal period Number of days in life cycle Number of generation each year	2 days 9-26 days 2-8 days 15-34 days 4-5	Ogah and Nwilene, 2017; Bhatt et al., 2018; Ouattara et al., 2020.

Table 2. Infestation parameters of rice pests

Rice pest	Name of pest	Most infectious stage	Economic threshold level	Nature of damage	Symptom of damage	Reference
Stem borer	Yellow stem borer <i>Scirphophaga incertulas</i>	Larva	5–10% damaged tillers	Due to larval boring and feeding activity inside leaf sheath resulting in yellowing of leaves.	Dead Heart (DH)	Bhatt et al., 2018; Pathak and Khan, 1994; Baladhiya et al., 2018; Asghar et al., 2009; Cork et al., 1998.
	Pink stem borer <i>Sesamia infernce</i>	Larva	1.9-3.0% WHs			
	Stripped stem borer <i>Chilo suppressalis</i>	Larva	2-8% hill or damage tillers	On severe condition folding of older rice leaves became drying off and turns whitish in colour.	White Head (WH)	
	Dark headed stem borer <i>Chilo polychrysus</i>	Larva	5% DHs			
	Leaf folder <i>Cnaphalocrocis medinalis</i>	Larva	3% folded leaves or 0.2-5 adult per hill	Scrapping the green mesophyll tissue affecting on photo-system II activity in photophosphorylation process resulting in scorching and drying of the leaves affecting in plant growth and reduce yield.	Leaf folding	Padmavathi et al., 2012; Pathak and Khan, 1994; Bhatt et al., 2018; Asghar et al., 2009.
Plant-hoppers	Brown planthopper <i>Nilaparvata lugens</i>	Nymphs and adults	3-8 nymphs/hill	Sucking phloem sap from stem and leaf sheath resulting yellowing of leaves and by ovipositional activity causing damage to plant tissue. On severe infestation results complete drying of the plants. Transmit Rice grassy stunt virus (RGSV) and rice ragged stunt virus (RRSV).	Hopperburn, Grassy Stunt (GS), Ragged Stunt (RS), Witte Stunt (WS), and Rice Yellowing Syndrome (RYS)	Catindig et al., 2009; Bhatt et al., 2018; Kaur et al., 2015; Bottrell and Schoenly, 2012; Matharu and Tanwar, 2020.
	White backed planthopper <i>Sogatella furcifera</i>	Nymphs and adults	4% hopperburn or 10-20 hoppers/hill	Sucking phloem sap from stem and leaf sheath resulting yellowing of leaves and by ovipositional activity causing damage to plant tissue and encourages the growth of sooty mould eg. <i>Cladosporium</i> sp and <i>Dematium</i> sp. On severe infestation results complete drying of the plants. Transmit Southern rice black-streaked dwarfvirus (SRBSDV).	Hopperburn, Southern rice black-streaked dwarf disease (SRBSDDD)	Catindig et al., 2009; Kaur et al., 2015; Bhatt et al., 2018.
	Green leafhopper <i>Nephotettix virescens</i> and <i>Nephotettix nigropictus</i>	Nymphs and adults	2-20 insects per hill	Suck up sap from vascular vessels and by ovipositional activity damage cortex and epidermal layer of leaf resulting yellowing plant, drying up of panicle. Transmit Rice Tungro Viruses (RTV), Rice Tungro Bacilliform Virus (RTBV), Rice Tungro Spherical Virus (RTSV), Rice Dwarf Virus (RDV), Rice Waika Virus (RWV) along with bacterial and fungal infection.	Milder Tungro Symptoms (MTS), Mild Stunting Symptoms (MSS), Dwarf Syndrome (DS), Yellow-Orange Leaf Symptom (YOLS)	Pathak and Khan, 1994; Dey, 2016; Sabale et al., 2010.

(contd.)

(Table 2 contd.)

Rice hispa	Rice hispa <i>Dictyodisca armigera</i>	Grubs and adults	2 damaged leaves/plant or 1-2 adults/hill	By feeding upon the chlorophyll content of leaves and also through scrapping the upper and lower surfaces of leaves resulting appearance of blisters and blotches or burned appearance on surface of leaves with parallel white lines on the leaf surface.	Leaf folding	Iqbal et al., 2015; Rahaman et al., 2020.
Rice hug	Gundhi bug <i>Leptocoris oratorius</i>	Nymph and adult	2 bugs/ m ² or 2 adults/ hill	Through piercing and sucking mouth part suck up the milky juice of the developing rice grains resulting partially filled grains. Severe infestation resulting chaffy and empty grains remain small in size. Passively transmit <i>Sarocladium oryzae</i> and <i>S. attenuatum</i> (fungi).	Panicles discoloration, Sooty Mould, Sheath Rot	Dutta and Roy, 2016; Khare and Prakash, 2020; Bhatt et al., 2018; Arora et al., 2019; Dutta et al., 2020, 2021; Bhatt et al., 2018.
	Earhead bug <i>Leptocorisa acuta</i>	Nymph and adult	0.5-1 individuals (adult and nymph)/ hill			
	Rice mealy bugs <i>Brevinnia rehi</i>	Nymphs and adults	3-5% affected tillers	Sucking the sap from stems and leaf sheaths resulting in leaves become yellowish, dried off and curled, also affects the panicle-bearing capacity of plant. Act as a vector for chlorotic streak virus and <i>Sarocladium oryzae</i>	Stunting and wiltingplant, Sheath Rot Disease	Bhatt et al., 2018; Pathak and Khan, 1994; Rath et al., 2020b.
Gall midge	Rice gall midge <i>Orseolia oryzae</i>	Larval	5-10% onion shoots or silver shoot	Feeding secretes a chemical known as cecidogen from its saliva which affecting nutrient supply and resulting central leaf elongation and tubular gall-like structure is formed. Larval feeding on the meristematic tissue of terminal and auxiliary shoot apices resulting in gall formation.	Onion leaf Silver shoot	Ogah and Nwilene, 2017; Bhatt et al., 2018; Kumari et al., 2020.

leaves longitudinally with the abaxial surface inside the folds, while younger instars partially rolled the leaf edges and folds were made along the middle portion of upper leaves with drying off central leaf which turn into whitish in colour and condition is known as 'White Ears or White Head' (WH) (Baladhiya et al., 2018). Agrosystem analysis estimated that economic threshold level was 5 to 10% damaged tillers for *Scirphophaga incertulas* (Walker) and 1.9-3.0% white heads for *Sesamia inferenze* (Walker) whereas 2-8% hill or damage tillers for *Chilo suppressalis* (Meyrick) while 5% dead hearts recorded for *Chilo polychrysus* (Meyrick) (Asghar et al., 2009; Cork et al., 1998). Stem borers activity mostly has been recorded mostly on rice, beside that they also aggregate on wild grasses like *Paniculum* sp., *Echinochloa* sp., *Cyperus* sp., *Dactylis* sp., *Scirpus* sp., *Saccharum* sp. etc. grown in and around crop fields (Bhatt et al., 2018; Dutta and Roy, 2018b; Jeevanandham et al., 2020).

B. Leaf folder

Rice leaf folder, *Cnaphalocrocis medinalis* (Guenee, 1854) (Lepidoptera: Pyralidae) were predominant defoliator pests during nursery to harvest (Bhatt et al., 2018). It encompasses egg, 5-6 larval instars, pupa and adult (Gangwar, 2015). Damage generally done due to voracious feeding by fourth and fifth instar inside leaf by scrapping the green mesophyll tissue of upper epidermis but the lower epidermis becomes dry and falls off and longitudinal folding of leaves by stitching the leaf margins with the help of salivary secretions (Padmavathi et al., 2012). Within the folded leaves reduction in chlorophyll content occur, influencing on photo-system II activity in photophosphorylation process in photosynthesis resulting membranous patches resulting in scorching and drying of the leaves which ultimately affecting in plant growth and reduce yield (Pathak and Khan, 1994). Beside that faecal matter of insect inside the folded leaf can cause development of fungal and bacterial infections (Bhatt et al., 2018). Researches have shown that economic threshold level which are 3% folded leaves or 0.2-5 adult per hill (Asghar et al., 2009). This deleterious leaf folder mainly infested rice with that they aggregated on plant like maize, *Panicum decompositum*, *Leersia hexandra*, *Echinochloa crusgalli*, *Leptochloa chinensis*, *Isachne dispar*, *Eleusine coracana*, and *Pennisetum typhoides* (Hurali et al., 2020; Asghar et al., 2009).

C. Planthopper

Herbivorous sap-feeding homopterans brown

plant hopper (BPH) *Nilaparvata lugens* (Stal, 1854) (Hemiptera: Delphacidae) and white backed plant hopper (WBPH) *Sogatella furcifera* (Horvath, 1899) (Hemiptera: Delphacidae) have posed a serious threat in summer Amon rice of Bengal (Matharu and Tanwar, 2020). Generally, two morphological types of plant hopper, macropterous and brachypterous have noticed in agrosystem (Catindig et al., 2009). Both morphological types of *N. lugens* (Stal) and *S. furcifera* (Horvath) encompass egg, six nymphal stages and adult (Bottrell and Schoenly, 2012). Damages caused by nymphs and adults of *N. lugens* (Stal.) and *S. furcifera* (Horvath) actively in growing plants during tillering stage and flowering stage by sucking phloem sap from stem and leaf sheath resulting yellowing of leaves and by ovipositional activity during booting and heading stages causing damage to plant tissue (Catindig et al., 2009; Bhatt et al., 2018). Severe infestation results complete drying of the plants, known as hopperburn (Kaur et al., 2015; Catindig et al., 2009). Feeding punctures and lacerations caused by ovipositor predispose of white-backed planthopper on plants to pathogenic organisms and honey dew excretion encourages the growth of sooty mould such as *Cladosporium* sp. and *Dematium* sp. (Kaur et al., 2015). Passively *N. lugens* (Stal.) can damage by transmitting Rice grassy stunt virus (RGSV) and rice ragged stunt virus (RRSV) result in rice yellowing syndrome (RYS) in rice (Bottrell and Schoenly, 2012). Similarly, *S. furcifera* (Horvath) act as an important vector of fiji virus, Southern rice black-streaked dwarf virus (SRBSDV) causing of Southern rice black-streaked dwarf disease (Bhatt et al., 2018). ETLs have encountered for brown planthopper 3-8 nymphs/hill (Matharu and Tanwar, 2020) whereas 4% hopperburn or 10-20 hoppers/ hill have recorded for white-backed planthopper (Bhatt et al., 2018). Generally, the rice is considering the best suitable host system for planthopper development but they had also aggregate on a weed *Leersia* sp., *Digitaria* sp., *Echinochloa* sp., *Isachne globosa*, *Poa annua*, *Poa repens* (Bhatt et al., 2018; Matharu and Tanwar, 2020).

D. Leafhopper

The rice green leafhoppers (GLHs) *Nephotettix virescens* (Distant, 1908) and *Nephotettix nigropictus* (Stal, 1870) (Hemiptera: Cicadellidae) have considered another devastating herbivorous pest of Kharif crop season (Bhatt et al., 2018). During life cycle it encompasses Egg, five instars nymph and adult stages (Dey, 2016). Damage mainly cause by either directly with their needle shaped piercing and sucking mouth

part sucking the sap and plugging the vascular bundles (xylem and phloem) with their feeding stylet sheaths results in wilting and pieces of tissue pushed into these vessels during exploratory feeding and oviposition or indirectly by transmitting virus diseases during feeding caused by Rice Tungro Viruses (RTV), Rice Tungro Bacilliform Virus (RTBV), Rice Tungro Spherical Virus (RTSV), Rice Waika Virus (RWV), Rice dwarf virus (RDV) responsible for Milder Tungro symptoms, Mild stunting Symptoms Wilting, Dwarf syndrome, Transitory yellowing syndrome, Yellow dwarf syndrome and Yellow-orange leaf symptom and also transmit fungal and bacterial infection responsible for Sooty Moulds in infected plant (Pathak and Khan, 1994; Dey, 2016). Economic threshold level of leafhopper has found 10 ± 9 insects per hill (Sabale et al., 2010). Generally, the rice is considering the best suitable host system for planthopper development but they had also aggregate on *Leersia* sp., *Digitaria* sp., *Echinochloa* sp., *Isachne globosa*, *Poa annua*, and *Panicum repens* etc. (Sarao and Bentur, 2018; Dey, 2016).

E. Rice hispa

The rice hispa or spiny beetle, *Dicladispa armigera* (Olivier, 1808) (Coleoptera: Chrysomelidae), has occupied a status of major pest of boro rice cultivation (Bhattacharjee and Ray, 2017). This pest encompasses egg, four instars larva or grubs, pupa and adult during their development (Iqbal et al., 2015). During vegetative stage of plant, the grubs feed upon the chlorophyll content of leaves and leaves became dry and give rise to blisters or blotches (Bhattacharjee and Ray, 2017). Adults also damaged through scrapping the upper and lower surfaces of leaves result in white streaks parallel to the midrib of leaves (Iqbal et al., 2015). During infestation photosynthetic activity of the leaves gets affected and leaves became dry and give rise blisters and blotches or burned appearance on surface of leaves with parallel white lines on the leaf surface (Rahaman et al., 2020). It has reported that ETL of this delictorious pest recorded 2 damaged leaves/plant or 1-2 adults/hill (Iqbal et al., 2015). This notorious pest also aggregated on *Andropogon gayanus*, *Sorghum halepense*, *Imperata cylindrica* and *Themeda anathera* (Rahaman et al., 2020).

F. Rice bug

The rice bugs have considered prevalent major pests for rainfed wetland or upland rice belong to *Leptocorisa* sp., *Brevennia rehi* (Lindinger, 1943) (Rath et al., 2020a, 2020b). *Leptocorisa acuta* (Thunberg, 1783) (Hemiptera: Alydidae) and *Leptocorisa*

oratorius (Fabricius, 1794) (Hemiptera: Alydidae) are known as Gundhi Bug (GB) due to releasing of characteristic graceless, bothersome, incursive odour on inciting caused by chemicals released from the bug's abdominal glands (Dutta et al., 2020). Rice Mealy Bugs (RMB), *Brevennia rehi* (Lindinger, 1943) (Hemiptera: Pseudococcidae) are plant-sucking, sporadic insect pest of rainfed rice (Bhatt et al., 2018). Life cycle of *Leptocorisa oratorius* (Fabricius) include egg, five nymphal stage and adult (Khare and Prakash, 2020; Arora et al., 2019) but *Leptocorisa acuta* (Thunberg) also shown developmental stages except they have six nymphal stages instead of five stages (Dutta and Roy, 2016, 2018a). *Brevennia rehi* (Lindinger, 1943) has encompassed egg, four nymphal instars and adult during their development (Bhatt et al., 2018; Pathak and Khan, 1994). Damage occurs generally during milk stage to dough stage by nymphs and adults of *L. acuta* (Thunberg) and *L. oratorius* (Fabricius) through piercing and sucking mouth part suck up the milky juice of the developing rice grains resulting partially filled grains (Dutta and Roy, 2016, Khare and Prakash, 2020; Dutta et al., 2020). Severe infestation resulting chaffy and empty grains remain small in size (Bhatt et al., 2018). The *L. acuta* (Thunberg) and *L. oratorius* (Fabricius) passively transmit *Sarocladium oryzae* and *S. attenuatum*, the cause of sheath rot disease which results in damaging of the panicle of the rice plant and infected plant may not produce rice grains (Arora et al., 2019). Economic threshold level has encountered for *L. acuta* (Thunberg) 2 bugs/ m² or 2 adults/ hill (Dutta et al., 2020) whereas that level for *L. oratorius* was 0.5-1 individuals (adult and nymph)/ hill (Bhatt et al., 2018). Similarly, damages occur by *B. rehi* (Lindinger) by both nymphs and adults feed by sucking the sap from stems and leaf sheaths resulting leaves become yellowish, dried off, and curled (Bhatt et al., 2018). The infestation also affects the panicle-bearing capacity of plant and produces chaffy grains and also stunting and wilting infected plant leading growth and development of the affected plant with that infested fields show isolated patches of stunted plants (Rath et al., 2020b). Presence of this bug well marked by protruding white waxy fluff between the leaf sheath and the stem and mealybug-infested grains have not developed properly, having bitter taste with spoil, flavour (Pathak and Khan, 1994). Passively this insect pest is also known to transmit chlorotic streak virus and *Sarocladium oryzae* responsible for sheath rot disease (Bhatt et al., 2018). Economic threshold level of *B. rehi* (Lindinger) has estimated 3-5% affected tillers (Rath et al., 2020b). *L. acuta* (Thunberg), *L. oratorius* (Fabricius) and *B. rehi*

(Lindinger) are generally aggregated on rice host (*Oryza sativa*) beside that they also aggregated on non rice hosts also like, *Echinochloa glabrescens*, *Paspalidium flavidum*, *Paspalum scrobiculatum*, *Echinochloa crusgalli*, *Echinochloa colona*, *Paniculum* spp. *Cyperus* sp. (Dutta and Roy, 2018a; Rath et al., 2020b).

G. Gall midge

Asian rice gall midge, *Orseolia oryzae* (Wood-Mason, 1889) (Cecidomyiidae: Diptera) were considered as one of the major destructive pests of lowland to upland rice (Nacro et al., 1996). Development of *O. oryzae* (Wood-Mason) involves egg, three larval instars, pupa and adult completed in 15-34 days (Ogah and Nwilene, 2017). Gravid females are laid tubular or rod-shaped eggs in vicinity of plant below or above ligules, on leaf blade, central whorl, standing water near the plant (Mathur and Rajamani, 1984). Infestation by *O. oryzae* (Wood-Mason) has seen from nursery to the tillering stage (Bhatt et al., 2018). Larvae enter the shoot apex and larval feeding secretes a chemical known as cecidogen from its saliva which affecting nutrient supply and resulting central leaf elongation and tubular gall-like structure is formed, resembles as “onion leaf” beside that due to larval feeding on the meristematic tissue of terminal and auxiliary shoot apices resulting in gall formation the characteristic symptom is known as “silver shoot” due to the shiny appearance of the leaf (Ogah and Nwilene, 2017). Through mathematical model that ETLs are 5-10% onion shoots or silver shoot (Bhatt et al., 2018; Kumari et al., 2020). Rice *Oryza sativa* has considered as most favourable host but this pest also aggregated in *Cynodon dactylon*, *Echinochloa colona*, *Paspalum scrobiculatum*, *Brachiaria mutica*, *Heteropogon contortus*, *Paspalidium geminatum* and *Leersia hexandra* (Bhatt et al., 2018).

H. Other pests

There are other insect pests of rice in West Bengal having regional significance such as army worm *Mythimna separate* (Walker), grasshoppers *Hieroglyphus banian* (Fabricius), white stem borer *Scirpophaga innotata* (Walker) and rice thrips *Stenchaetothrips biformis* (Bagnall), etc. (Chatterjee et al., 2008).

Management strategies (Table 3)

In this modern era with increasing human population there is a need to increase rice production per unit of land through sustainable management strategies (Dutta et al., 2021; Chávez et al., 2018). To minimize

yield losses and to increased productivity farmers use different management strategies with minimum cost to control pest population in agro ecosystem as described below (Roy, 2019a, 2019b; Kunal et al., 2020).

A. Chemical control

Most popular method used by farmers for controlling insect rice pest was through using synthetic chemical pesticide (Roy, 2019a, 2019b). Application of aqueous or oily spray of fenvalrate, acephate, quinalphos, cartap hydrochloride, flubendiamide, carbofuran, carbosulfan, chlorantraniliprole, fipronil, thiamethoxam, chlorpyrifos, spinosad, and N-((2,6-dimethyl-4-(heptafluoropropyl-2-yl)phenyl) carbamoyl)-2,6-difluorobenzamide has shown best efficacy against adult and larva of rice leaf folder, gall midge, planthopper, leafhopper, gandhi bug, mealybug and stem borers (Shyamrao and Raghuraman, 2019; Dutta and Roy, 2022; Zhao et al., 2019; Baladhiya et al., 2018). Application of soil granular formulation of diazinon, propaphos, endrin chlorfenvinphos, disulfoton, padan, deltamethrin, propenphos, propenphos, chlopyriphos, monocrotophos, imidacloprid, buprofezin, cycloxaprid, pymetrozine, sulfoxaflor, indoxacarb, flonicamid, malathion, dimethoate, phosphamidon, dinotefurain, chlorpyriphos, cypermethrin, lindane, triazophos, spinosad, rynaxypyr, and endosulfan during seedling plantation have shown good result for managing adult and larval incidence caused by major insect pests (Dutta and Roy, 2022; Zhao et al., 2019; Achiri et al., 2020; Matharu and Tanwar, 2020; Hurali et al., 2020; Luo et al., 2023; Adhikari et al., 2019; Bhattacharjee and Ray, 2017; Gupta et al., 2019; Kumari et al., 2020; Adhikari et al., 2019). Monocrotophos, imidacloprid, and triazophos were most economical on basis of cost benefit ratio (CBR) (Tigga et al., 2018; Adhikari et al., 2019).

B. Cultural control

For controlling deleterious rice pests, popular mechanical and cultural techniques used by farmers was based on crop sanitation because in most of cases it has observed that source of infestation occurred by egg, larva and adult of phytophagous rice insect pests that persist during seedlings (Ayelo et al., 2021). Another popular method was based on plant quarantine technique which includes raising crop nurseries away from light source avoids the early infestation have done by rice insect pests (Seleiman et al., 2021). Most commonly used crop sanitation technique are seed soaking with fertilizer along with insecticide, clipping

Table 3. Management strategies for rice pests

Rice pest	Common name and scientific name	Management agents	Management strategies	References
		Carbofuran, carbosulfan, fipronil, acephate, fipronil, chlorpyrifos, quinalphos, flubendiamide.	Chemical	Shyamrao and Raghuraman, 2019; Dutta and Roy, 2022
	Yellow stem borer <i>Scirphophaga incertulas</i>	Leaf extract of neem (<i>Azadirachta indica</i>), tobacco (<i>Nicotina tobacum</i>), Karanja (<i>Pongamia glabra</i>), bishkatali (<i>Polygonum hydropiper</i>), lantana (<i>Lantana camara</i>) and neem oil, mahogany (<i>Swietenia macrophylla</i>) oil, etc.	Botanicals	Shyamrao and Raghuraman, 2019.
		Extract of <i>Trichogramma japonicum</i> .	Biological	Dutta and Roy, 2022.
	Pink stem borer <i>Sesamia inferens</i>	Fenvalrate, thiamethoxam, spinosad, chlorantraniliprole, carbofuran.	Chemical	Baladhya et al., 2018; Zhao et al., 2019.
		Isolates of earwig <i>Euborellia stali</i> (Dohrn), Bt protoxin CryIca, Isolates of <i>Beauveria bassiana</i> , NBAlI- Bb-1, 4, 6, 8, 11, 15, 39 and 59 isolates.	Biological	Baladhya et al., 2018.
	Dark headed stem borer <i>Chilo suppressalis</i>	Fipronil, carbofuran, diazinon, disulfoton Padan, TXH09 [N-((2,6-dimethyl-4-(heptafluoropropyl-2-yl)phenyl)carbamoyl)-2,6-difluorobenzamide]	Chemical	Zhao et al., 2019; Khan et al., 2020.
		Extract of rice variety TKM-6, Pentadecanol TKM-6	Botanicals	Prakash et al., 2008.
		Extract powder of <i>Trichogramma japonicum</i> , <i>Trichogramma chilonis</i> , <i>T. dendrolimi</i> , and <i>T. ostriniae</i>	Biological	Hajjar et al., 2023.
	Stripped stem borer <i>Chilo polychrysus</i>	Spinosad, deltamethrin, lambda-cyhalothrin, emamectin-benzoate, indoxacarb, chlorpyrifos-ethyl, chlorantraniliprole, novaluron	Chemical	Achiri et al., 2020.
		Extract of <i>Neorautanenia mitis</i> , <i>Derris elliptica</i> , <i>Metarhizium anisopliae</i> , <i>Beauveria bassiana</i>	Biological	January et al., 2020.
		Fipronil, propenphos, chlorpyrifos, monocrotophos, carbofuran, carbosulfan, imidacloprid	Chemical	Hurali et al., 2020.
	Leaf folder <i>Cnaphalocrocis medinalis</i>	Azadirachtin, salannin, deacetylgedunin, gedunin, 17-hydroxyazadiradione, deacetylhinmbin	Botanicals	Nathan et al., 2005.
		Pseudomonas stain (Pfl + TDK1 + PY15), Cnme GV, (isolated from <i>Yangzhou</i>)	Biological	Jian et al., 2019.
		Endrin, carbofuran, disulfoton, chlorfenvinphos, phorate, methyl-demeton, dicrotophos, monocrotophos, imidacloprid, flonicamid, etc.	Chemical	Matharu and Ianwar, 2020.
		Extract of <i>Adhatoda vasica</i> , <i>Vitex negundo</i> , <i>Azadiracta indica</i> , <i>Ricinus comucies</i> and <i>Pongamia glabra</i> leaves, neem seed kernal extract, etc.	Botanicals	Longkumer and Misra, 2020.
	Brown planthopper <i>Nilaparvata lugens</i>	Conidial suspension of <i>Metarhizium anisopliae</i> , <i>M. flavoviride</i> , <i>Beauveria bassiana</i> , and <i>Hirsutellatritiformis</i>	Biological	Sharma and Sharma, 2021
		Imidacloprid, buprofezin, thiamethoxam, chlorpyrifos, isoprocarb, cycloxaprid, buprofezin	Chemical	Luo et al., 2023.
	White backed planthopper <i>Sogatella furcifera</i>	Leaf crude extract of neem, notchi (<i>Vitex negundo</i>), palmarosa (<i>Cymbopogon mauritii</i>) oil, jatropa (<i>Jatropha carcus</i>) oil, neem oil, etc.	Botanicals	Raj et al., 2020.
		Jinggangmycin (JGM) [product of <i>Streptomyces</i> var. <i>jinggangensis</i>]	Biological	Ge et al., 2017.

(contd.)

(Table 3 contd.)

Leafhopper	Green leafhopper <i>Nephotettix virescens</i> and <i>Nephotettix nigropictus</i>	Pymetrozine, imidacloprid, dinotefuran, sulfoxaflor, flonicamid, buprofezin, propaphos, disulfoton	Chemical	Adhikari et al., 2019.
		Karanj oil, mahua oil, pinnai oil, neem oil, custard-apple oil, seed oil and leaf and fruit extract of <i>Annona squamosa</i>	Botanicals	Kumar et al., 2021.
		Conidia and blastospores of <i>Beauveria bassiana</i>	Biological	Sharma and Sharma, 2021.
		Quinalphos, malathion, dimethoate, phosphamidon, endosulfan, chlorpyrifos, monocrotophos, cypermethrin, fenvalarate	Chemical	Bhattacharjee and Ray, 2017.
Rice hispa	Rice hispa <i>Diuraphis armigera</i>	Neem oil, Azacel, Calpaste, Larvoce, dk-bioneem, Multineem Powder formulation of <i>Beauveria bassiana</i> (Bals.) Vuill, spore extraction of <i>B. bassiana</i>	Botanicals	Kumar et al., 2021.
		Imidacloprid, triazophos, monocrotophos, thiamethoxam, acephate, carbaryl, malathion, lindane, etc.	Biological	Purnima et al., 2013.
		Extract of <i>Chromolaena odorata</i> , <i>Azadirachta indica</i> , <i>Ricinus communis</i> , neem seed kernel powder (NSKP), etc.	Chemical	Gupta et al., 2019.
Rice bug	Earhead bug <i>Leptocorisa acuta</i> , gundhi bug <i>Leptocorisa oratorius</i> , rice mealy bug <i>Brevinnia rehi</i>	Isolates of entomopathogens (<i>Beauveria bassiana</i> and <i>Metarhizium anisopliae</i>), cow urine	Botanicals	Raveau et al., 2020.
		Acephate, rynaxypyr, dinotefuran, fipronil, thiamethoxam, buprofezin, spinetoram, methoxyfenozide, acephate, dinotefuran, triazophos, Neem Seed Kernel Extract (NSKE)	Biological	Kumar and Sarada, 2020.
Gall midge	Rice gall midge <i>Orseolia oryzae</i>	Parasitized galls of <i>Platygaster diplosisae</i>	Chemical	Kumari et al., 2020.
			Botanicals	Ogah and Nwile, 2017.
			Biological	Minab et al., 2023.

of leaves of rice plants during highest tillering stage, removal of stubbles through plough and then burn at ground, drain out water in infested paddy field, seedling root dipping, removal of alternate grassy and other weeds hosts from seedbed, crop field and adjacent area shown very effective against leaf folder, stem borer, gall midge, rice hispa and rice bug (Bhatt et al., 2018; Kunal et al., 2020). Cultivation of certain susceptible aromatic rice cultivars like Pusa Basmati-1, Gobindabhog near main crop can be used as trap crop to reduce damage to the main crop (Katti et al., 2023). Crop rotation with leguminous pulse seed, potato, oil seed in low land and cultivation of maize in upland area are also shown good efficacy against different insect pests (Breidenbach et al., 2017; Chen et al., 2018). Uses of oil or electric light traps are very helpful against stem borer and planthopper (Bhatt et al., 2018). Even, use of fish like, *Puntius conchoni* (F. Hamilton, 1822), *Clarias batrachus* (Linnaeus, 1758), *Punctius ticto* (F. Hamilton, 1822), frogs like, *Euphyctis cyanophlyctis*, *Duttaphrynus melanostictus* (Schneider, 1799) etc. and *Anas poecilorhyncha* (Forster, 1781) are preyed on adult of rice pest thereby reduce the population (Khatiwada et al., 2016; Yuan et al., 2022).

C. Biopesticides

Plant formulations, so called 'green pesticide' were also effective and have minimal residual activity (Raveau et al., 2020). Natural oil, cake and crude extract of different part of neem plant [*Azadirachta indica* (Juss, 1830)] neem oil, neem oil soap solution, neem seed oil, cake extract, leaf extract, leaf crude extract, seed kernel extract, limonoid, salannin, deacetylgedunin, gedunin, 17-hydroxyazadiradione, deacetyl nimbin, azadirachtin, neemgold, nimbidine, neemax, azacel, and larvocel have shown good efficacy against deleterious pest incidence (January et al., 2020; Shyamrao and Raghuraman, 2019; Dougoud et al., 2019). Application of extract of different part from other plants like seeds of *Pongamia pinnata*, *Madhuca longifolia*, *Calophyllum inophyllum*, *Neorautanenia mitis*, *Derris elliptica*, *Nicotinatabacum*, *Pongamia glabra*, *Chromolaena odorata*, *Ricinus communis*, *Adhatoda vasica*, *Andrograpis paniculata*, *Prosopis juliflora*, *Polygonum hydropiper*, *Lantana camara*, *Vitex negundo*, mixture of *Polygonum hydropiper*, *L. camara* and ginger-garlic-chili effectively reduced incidence caused by herbivores insect pests (Raj et al., 2020; Longkumer and Misra, 2020). Oil formulation from Mahogany, Palmarosa, Jatropha, custard-apple seed, Karanj (*Pongamia pinnata*), Eucalyptus (*Eucalyptus*

obliqua), Cedar Wood, Mahua (*Madhuca longifolia*), Pinnai (*Calophyllum inophyllum*), Citronella, Vetiver, Catnip, were effective against *Di cladispa armigera* (Olivier), *Sogatella furcifera* (Horvath), *Nephotettix virescens* (Distant), *Nephotettix nigropictus* (Stal), *Scirphophaga incertulas* (Walker) population (Raj et al., 2020; Kumar et al., 2021; Raveau et al., 2020). Animal derived cow urine also acts as antifeedant and disinfectant against rice pests and their transmitting pathogens (Kumar and Sarada, 2020).

D. Biological control

Most popular biological controlling agents include mycobial powder and pesticide extracted from *Bacillus thuringiensis*, *Trichogramma japonicum*, *T. chilonis*, *T. dendrolimi*, *T. ostrinae*, *Lecanicillium lecanii*, *Beauveria bassiana*, *Metarhizium anisopliae*, *Skermanella* sp., *Pseudomonas* stain, CnmeGV galls of *Platygaster diplosisa* and Jingga mycin (JGM) had shown good antibiosis effects through egg parasitic, larvicidal activity on rice insect pests (Dutta and Roy, 2022; Hajjar et al., 2023; January et al., 2020; Kumar and Sarada, 2020; Minab et al., 2023; Jian et al., 2019; Ge et al., 2017). Using of conidial suspension of *Metarhizium anisopliae*, *M. flavoviride*, *Beauveria bassiana*, and *Hirsutella citrififormis*, preparations of marcescent mycelium of *Paecilomyces lilacinus*, wettable powder formulation, spore extract, conidia and blastospores extract of *Beauveria bassiana* had shown egg parasitoid activity on *C. polychrysus* (Meyrick), *D. armigera* (Olivier), plant and leafhopper (Purnima et al., 2013; Sharma and Sharma, 2021). Bio-pesticides Btprotoxin CryIca, NBAII-Bb-1, 4, 6, 8, 11, 15, 39 and 59 isolates, NBAII-Ma-1, 6, 10, 13, 15, 23, 25, 26, 41 and 42 isolates effective against *D. armigera* (Olivier) population (Baladhiya et al., 2018).

E. Resistant variety based control

Rice variety D518, IR-42, ASD7, and IR46 were effective against incidence by *N. lugens* (Stal.) and *S. furcifera* (Horvath) (Gao et al., 2020). Infestation potential of *C. medinalis* (Guenee) has recorded low in cultivar Ptb-12, TKM-6, Prasanna, Radha-4, Ramdhan, IRRI-6, KSK-282, DR-83, Bas-385 (Adhikari et al., 2022; Chen et al., 2022; Chatterjee et al., 2021). Similarly, GAR-13, IR-22, Gontra Bidhan 3, MTU-2020, IR-50, TKM 6, KMD1, KMD2, MTU15, and IR50 have shown good efficacy against *S. incertulas* (Walker), *C. suppressalis* (Meyrick) and *C. polychrysus* (Meyrick) (Jha et al., 2023; Chavan and Patel, 2018a, 2018b; Xu et al., 2022). Pankhari 203, WRC25,

WRC34, WRC37, WRC40, WRC6, WRC7, WRC48, and WRC55 has recorded effective against *N. virescens* (Distant) and *N. nigropictus* (Stal) (Thein et al., 2019). Rice variety like IET-6286, Surekha, Usha, Sahbhagi Dhan, Sudha, Abhisek, ARC-5833, ARC-5984, TKM-6, and PTB-10 were effective against *O. oryzae* (Wood-Mason) and *S. inference* (Walker) incidence (Sanchit and Sanchit, 2019). Rice cultivar like Garem and Bengaubisi found effective against *D. armigera* (Olivier) infestation (Dutta and Hazarika, 1994). GR-9, Jaya, Dandi, Pawana, Gurjari, Narmada, Pawana, and GR-104 have shown good efficacy against *B. rehi* (Lindinger), *L. acuta* (Thunberg) and *L. oratorius* (Fabricius) (Khalifa and El-Rewainy, 2012; Morya and Kumar, 2019; Rath et al., 2020b).

F. Phoromone and allomone based control

Now a days farmers often use attractant sex pheromones (Z)-11-hexadecinal, (Z)-9-hexadecenal, (Z)-9-hexadecenal and (Z)-11-hexadecenal with PVC matrix, Z11-16 Acetate, (Z)-hexadec-9-enal, (Z)-11-octadecenal, (Z)-11-hexadecenyl acetate, (Z)-13-octadecenyl acetate, pymetrozine, 2-(E)-octenyl acetate, octanol, n-tridecane and 2S,6S-diaxetoxyheptane which help in attract male and female mass trapping and mating disruption of insects for management of rice pest (Gunawardena and Ranatunga, 1989; Dutta and Roy, 2022; Liu et al., 2020; Dang et al., 2016; Song et al., 2022). Some repellent allomone 5: 1 mixture of 2-(E)-octenyl acetate and octanol 3-octenal, 1-octanol, (Z)-3-octenyl acetate, extracts of Bamboo (*Bambusa levis* L.), Ginger (*Catimbium haenkei* L.), Malubago (*Hibiscus tiliaceus* L.) repel herbivore stem borer, leaf folder and gandhi bug population (Gunawardena and Bandumathie, 1993).

G. Population based control

Several demographical models described the biotic and abiotic mortality factors affecting its abundance, development, survivorship, reproduction, in agroecosystem (Mobarak et al., 2020). Developing of age-specific (horizontal) and stage-specific (vertical) life tables for assessing population density, fluctuation, most vulnerable stage or stage having higher mortality rate of different rice pests under different agroclimatic field conditions (Dutta and Roy, 2016; Dutta et al., 2020, 2021). Population growth and reproductive parameters [probability of survival from birth to age x (l_x), survival rate (s_x), average population alive in each stage (L_x), total individuals at age x and beyond k (T_x), life expectancy (e_x), generation survival

(GS), age specific fertility or fecundity (m_x), gross reproductive rate (GRR), net reproductive rate (NRR or R_0), intrinsic rate of population increase (r_m), finite rate of population increase (λ), weekly multiplication rate (λ^7), increase rate per generation (λ^{T_c}), total fertility rate (F_x), population growth rate (PGR), and population momentum factor of increase (PMF)], mortality profiles [proportion dying each age (d_x), mortality rate (q_x), and mortality coefficient (MC)], span of generation time (T_c) and doubling time (DT) depend on temperature, nutritional quality [quantity of carbohydrates, protein, lipid, phenol, phytate, saponin etc.] host system [rice host (*O. sativa*), non rice host (eg. *E. colona*, *C. rotundus*, etc.)], presence of egg and adult predator parasitoid (Babamir-Satehi et al., 2022; Schoonhoven et al., 2005; Dutta and Roy, 2018a, 2018b; Harborne, 1973, 1984; Manikandan et al., 2015; Havelka and Zemek, 1999). By analyzing life table data of rice pests in different host and agro-climatic condition type -II survivorship has shown by *S. furcifera* (Horvath) but type-III survivorship obtained in *S. incertulas* (Walker), *N. lugens* (Stal), *L. acuta* (Thunberg) and even, type IV exhibited by *C. suppressalis* (Meyrick) and *C. polychrysus* (Meyrick) (Dutta and Roy, 2016, 2018b; Shivasharanappa et al., 2010). By studying population dynamics, sixth nymphal stage of *L. acuta* (Thunberg), first instar larva of *S. incertulas* (Walker), *C. suppressalis* (Meyrick), *D. armigera* (Olivier), fourth nymphal stage of *N. lugens* (Stal) and first nymphal stage and adult of green leafhopper have identified most vulnerable stages (Win et al., 2009, 2011). Ecological limiting factors like relative humidity in winter, low moisture, predator density and dispersal loss during larval stage, egg predator [eg. *Conocephalus chinensis* (Redtenbacher)], high temperature, low relative humidity, parasitisation by the entomopathogenic nematode, adult migration, parasitoids of of egg, nymphal and adult, moulting failure of larva into pupa and host nutritional quality appeared as limiting factors *N. virescens* (Distant), *N. nigropictus* (Stal), *S. furcifera* (Horvath), *D. armigera* (Olivier), *C. suppressalis* (Meyrick), *L. acuta* (Thunberg) and *O. oryzae* (Wood-Mason) (Kumar et al., 2012; Dutta and Roy, 2016; Win et al., 2011; Suzuki et al., 1996; Roy, 2022). Parasitoids and disease on 3rd and 4th instar larva and pupal parasitism also appeared as limiting factor for *C. medinalis* (Guenee) (Huang et al., 2020; Padmavathi et al., 2008). By identifying most vulnerable stages and population dynamics along with limiting factors of rice insect pest are very essential for the timely adoption of different management practices (Mobarak et al., 2020; Roy, 2019a, 2019b, 2020, 2021).

CONCLUSIONS

Rice, one of the most consumable staple crop of most of Bengal household cultivated 56.63 lakh hectare areas with a production of 274.37 lakh tones, contributing about 20.69% to the total net State Domestic Product (SDP) (Dutta and Roy, 2022; Jha et al., 2023; Adak et al., 2020; Chakraborty and Deb, 2012). Rice in West Bengal has cultivated in the six agro-climatic zone during three different cropping seasons, Aus (autumn rice), Aman (winter rice) and Boro (summer rice) (Adhikari et al., 2012; Chakraborty and Deb, 2012). But this rice production in West Bengal was hampered by different biotic stresses causing 33% production loss (Aryal et al., 2018; Jha et al., 2023). Most important biotic stress caused to hamper production of rice in West Bengal was through direct or indirect infestation by phytophagous rice pest (Adhikari et al., 2012; Dutta and Roy, 2022). The remarkable deleterious rice insect pest fauna has observed mostly are *S. incertulas* (Walker), *S. inferens* (Walker), *C. polychrysus* (Meyrick), *C. suppressalis* (Meyrick), *C. medinalis* (Guenee), *N. lugens* (Stal.), *S. furcifera* (Horvath), *N. virescens* (Distant), *N. nigropictus* (Stal), *D. armigera* (Olivier), *L. acuta* (Thunberg), *L. oratorius* (Fabricius), *O. oryzae* (Wood-Mason) etc. (Azgar and Hembram, 2018; Dhaliwal et al., 2010; Jha et al., 2023).

It has observed that most of polyphagous major rice pests in rice mainly beside that they were also aggregated on some alternative non rice weeds found in and around rice field (Bhatt et al., 2018; Dutta and Roy, 2018a, 2018b). Population growth, reproductive and mortality parameters of different rice pests are greatly varied depending on host system and agroclimatic conditions (Roy, 2017, 2022; Reji and Chander, 2008). Modern agriculture concerned with ecofriendly and sustainable climate smart agriculture (CSA) as a part of ecological pest management (EPM) based integrated pest management (IPM) for of different economic crops (Pedigo et al., 1986; Pedigo and Higley, 1992; Aryal et al., 2018; Chávez et al., 2018; Roy, 2019b). Therefore, basic information regarding pest ecology is necessary before deciding any strategy to combat with deleterious rice pest (Rahaman et al., 2014; Roy, 2019b). Commonly most of farmers use different synthetic pesticides for controlling pests (Aktar et al., 2009; Bais et al., 2017). But some plant extract botanicals are also seen against rice pests (Aryal et al., 2018; Shitiri et al., 2014; Dutta and Roy, 2022). Beside these most popular controlling methods usually mechanical and cultural control strategies include mechanical synchronised

planting, cultivation of certain rice varieties like Pusa Basmati-1 as trap crop, soaking sprouted seed in synthetic pesticide, seedling root dipping in pesticide and fertilizer, remove, burn or plough the stubbles, and using of fish, frog ducklings etc. also seem to be useful for controlling these pest population (Ayelo et al., 2021; Khatiwada et al., 2016; Bhatt et al., 2018). Some attractant sex pheromones like, (Z)-11 hexadecinal, (Z)-9-hexadecenal, (Z)-11-hexadecenal, (Z)-9-octadecenal are show good efficacy in mass trapping and mating disruption of deleterious insect pests (Liu et al., 2020; Dang et al., 2016; Gunawardena and Ranatunga, 1989). Beside that repellent allomones like, 5: 1 mixture of 2-(E)-octenyl acetate and octanol 3-octenal, 1-octanol and extracts of resistant rice varieties distracts rice pests' population from cultivated field (Gunawardena and Bandumathie, 1993; Pathak and Khan, 1994). Farmers some time use biological agents as pesticides derived from strains of *Bacillus thuringiensis*, *Beauveria bassiana*, *Metarhizium anisopliae*, *M. flavoviride* etc. which were effective against major paddy pests (Dutta and Roy, 2022; Purnima et al., 2013; Sharma and Sharma, 2021). Certain resistant rice cultivars like, Pankaj, Triveni, IR-19362-183, IR-20, H4, IR-36, IR-50, IR-9828-23-1, IR-13429-57-1, IR-44, IR-22, and GR-9 etc. have impart resistance effect against rice pest population growth, development and reproductive parameters (Sanchit and Sanchit, 2019; Huang et al., 2019; Rath et al., 2020b).

Recent days, farmers generally use synthetic chemical insecticides indiscriminately against those pests for the hope of high production of rice, but it has seen that these pesticide not only target phytophagous rice pests but also targeted beneficial natural enemies of pest, pollinator resulting destabilizing of biodiversity, decrease the sustainability by bio-magnification and bio-accumulation of toxic substance which result in ecologically and environmentally imbalance and creates health hazards (Katti et al., 2023; Biswas et al., 2006; Aktar et al., 2009; Kumari et al., 2020). It has seen that excessive use of insect lethal insecticides leads to increase the amount of primary nutrient like amino acids and sucrose available in the phloem of rice plants results in insecticide-induced resurgence, and cross resistance leads to of periodic outbreaks pest populations, known as "Moran effect" (Bottrell and Schoenly, 2012; Dutta and Roy, 2016, 2022). To overcome this constrains caused by rice pest researchers usually use insecticide by considering several parameters like cost: benefit ratio, economic threshold level (Selvaraj et al., 2012), population dynamics by exploring of allelopathy-based

strategies and enhancing use of green pesticides and cultural methods for appropriate management strategies for rice pests (Dutta and Roy, 2022). Thus, population ecology based sustainable management of such pest species will support E³ strategy [Ecosystem service-based Ecological engineering for Ecological pest management (ESS-EE-EPM)] of pest management for successful cultivation of rice in the near future.

REFERENCES

- Achiri T D, Atakan E, Pehlivan S. 2020. Susceptibility of *Chilo partellus* Swinhoe, 1885 (Lepidoptera: Crambidae) to some commonly used insecticides. Turkish Journal of Entomology 44(2): 135-147. <https://doi.org/10.16970/entoted.609195>.
- Adak S, Datta S, Bhattacharya S, Ghose T K, Majumder A L. 2020. Diversity analysis of selected rice landraces from West Bengal and their linked molecular markers for salinity tolerance. Physiology and Molecular Biology of Plants 26: 669-682.
- Adhikari B, Bag M K, Bhowmick M K, Kundu C. 2012. Rice in West Bengal-rice knowledge management portal (www.rkmp.co.in). Directorate of Rice Research, Rajendranagar, Hyderabad, Andhra Pradesh. pp. 1-88.
- Adhikari B, Mohapatra L N, Senapati R, Mohapatra M, Muduli L, Mohapatra S D. 2022. Biochemical changes in rice leaves due to rice leaf folder *Cnaphalocrocis medinalis* (Guenee) infestation. The Pharma Innovation SP11(8).1463-1468.
- Adhikari B, Padhi J, Shitapragyan S. 2019. Efficacy of new molecules against green leafhopper in rice. Journal of Entomology and Zoology Studies 7(2): 194-197.
- Aktar M W, Sengupta D, Chowdhury A. 2009. Impact of pesticides use in agriculture: their benefits and hazards. Interdisciplinary Toxicology 2(1): 1-12.
- Al-Haddad, A S, mezel AL-Rubaei H, Dewan M M. 2018. Life cycle of rice Mealybug *Brevienia rehi* (Lindinger, 1943) in cultivars rice plants under the laboratory conditions. Plant Archives. 18 (1): 847-851.
- Arora S, Sehgal M, Srivastava D S, Arora S, Sarkar S K. 2019. Rice pest management with reduced risk pesticides in India. Environmental Monitoring and Assessment 191(241): 1-20.
- Aryal J P, Jat M L, Sapkota T B, Chhetri A K, Kassie M, Rahut D B, Maharjan S. 2018. Adoption of multiple climate smart agricultural practices in the Gangetic plains of Bihar, India. International Journal of Climate Change Strategies and Management 10(3): 407-427.
- Asghar M, Suhail A, Afzal M, Khan M A. 2009. Determination of economics threshold levels for the stem borers (*Scirpophaga* sp.) and leaf folder (*Cnaphalocrocis medinalis*) of rice (*Oryza sativa*) in the kallar tract of Punjab, Pakistan. International Journal of Agriculture and Biology 11: 717-720.
- Ayelo P M, Pirk C W, Yusuf A A, Chailleux A, Mohamed S A, Deletre E. 2021. Exploring the kairomone-based foraging behaviour of natural enemies to enhance biological control: A review. Frontiers in Ecology and Evolution, 9: 1-22.
- Azgar MA, Hembram K T. 2018. A comparative studies on the population dynamic and survival of green leaf hopper and zigzag leaf hopper transmitting rice tungro virus in West Bengal. The Bioscience 13(2): 607-611.
- Babamir-Satehi A, Habibpour B, Aghdam H R, Hemmati S A. 2022. Interaction between feeding efficiency and digestive physiology of the pink stem borer, *Sesamia cretica* Lederer (Lepidoptera: Noctuidae), and biochemical compounds of different sugarcane cultivars. Arthropod-Plant Interactions, 16(3-4): 309-316.
- Bag M, Adhikari B, Bhowmick M, Kundu C. 2011. Status paper on rice in West Bengal. Rice Research Station, Chinsura. Government of West Bengal, West Bengal. pp. 1-89.
- Bais B, Kumar A, Khan H H, Sahu R, Kaiwart P K. 2017. Efficacy of certain chemical insecticides against rice gundhi bug (*Leptocorisa acuta* Thunberg) under field condition. International Journal of Current Microbiology and Applied Sciences 6(9): 1697-1701.
- Baladhiya H C, Sisodiya D B, Pathan N P. 2018. A review on pink stem borer, *Sesamia inferens* Walker: a threat to cereals. Journal of Entomology and Zoology Studies 6(3): 1235-1239.
- Bhatt N, Joshi S, Tiwari S N. 2018. Pests of rice. In: Omkar (eds) Pests and their management, Springer, Singapore, pp. 9-50. https://doi.org/10.1007/978-981-10-8687-8_2.
- Bhattacharjee P P, Ray C D. 2017. Effect of season on the development of rice hispa, *Di cladispa armigera* in Barak Valley of Assam. Tropical Ecology 58(1): 199-203.
- Biswas T K, Rahman M A, Khan M M H, Alam M M, Jahan M. 2006. Toxic effects of two insecticides on brown plant hopper, *Nilaparvata lugens* and its predators *Micraspis discolor* and *Lycosa pseudoannulata*. International Journal of Zoological Research 2: 192-203.
- Bottrell D G, Schoenly K G. 2012. Resurrecting the ghost of green revolutions past: the brown planthopper as a recurring threat to high-yielding rice production in tropical Asia. Journal of Asia-Pacific Entomology 15(1): 122-140.
- Breidenbach B, Brenzinger K, Brandt F B, Blaser M B, Conrad R. 2017. The effect of crop rotation between wetland rice and upland maize on the microbial communities associated with roots. Plant and Soil 419: 435-445.
- Bunawan H, Dusik L, Bunawan S N, Amin N M. 2014. Rice tungro disease: from identification to disease control. World Applied Sciences Journal 31(6): 1221-1226.
- Carvalho F P. 2017. Pesticides, environment, and food safety. Food and Energy Security 6(2): 48-60.
- Catindig J L A, Arida G S, Baehaki S E, Bentur J S, Cuong L Q, Norowi M, Rattanakarn W, Sriratanasak W, Xia J, Lu Z, Heong K L. 2009. Situation of planthoppers in Asia. In (eds): Planthoppers: new threats to the sustainability of intensive rice production systems in Asia. International Rice Research Institute, Los Baños, Philippines. pp. 191-220.
- Chakraborty K. 2012. Effective management of *Scirpophaga incertulas* Walker on rice crop during kharif season in West Bengal, India. American-Eurasian Journal of Agricultural and Environmental Science 12(9): 1176-1184.
- Chakraborty K, Deb D C. 2012. Incidence of rice Hispa, *Di cladispa armigera* (Coleoptera: Crysomellidae) on kharif paddy in the agroclimatic conditions of the northern parts of Bengal, India. Global Journal of Science Frontier Research Biological Sciences 12(8): 52-61.
- Chatterjee S D, Adhikari B, Ghosh A, Ahmed J, Neogi S B, Pandey N. 2008. The rice bio-diversity in West Bengal. Department of Agriculture, Govt. of West Bengal. pp. 1-50.
- Chatterjee S, Gangopadhyay C, Dana I, Roy S K, Mondal P. 2021. Host plant resistance against yellow stem borer and rice leaf folder through varietal screening of rice. Plants and Environment 3(1): 23-29.
- Chavan S M, Patel K G. 2018a. Biochemical basis of resistance in rice

- varieties against yellow stem borer *Scirpophaga incertulas*. Indian Journal of Entomology 80(3): 1074-1079.
- Chávez J P, Jungmann D, Siegmund S. 2018. A comparative study of integrated pest management strategies based on impulsive control. Journal of Biological Dynamics Journal of Biological Dynamics 12(1): 318-341.
- Chen P, Dai C, Liu H, Hou M. 2022. Identification of key headspace volatile compounds signaling preference for rice over corn in adult females of the rice leaf folder *Cnaphalocrocis medinalis*. Journal of Agricultural and Food Chemistry 70(32): 9826-9833.
- Chen S, Liu S, Zheng X, Yin M, Chu G, Xu C, Yan J, Chen L, Wang D, Zhang X. 2018. Effect of various crop rotations on rice yield and nitrogen use efficiency in paddy-upland systems in south-eastern China. The Crop Journal 6(6): 576-588.
- Cork A, De Souza K, Krishnaiah K, Reddy AA, Zainullabuddin S. 1998. Season-long control of yellow stem borer, *Scirpophaga incertulas* (Lepidoptera: Pyralidae) by mating disruption with the natural ratio of pheromone components. Bulletin of Entomological Research 88(2): 109-116.
- Dang C H, Nguyen C H, Im C, Nguyen T D. 2016. Synthesis and Application of Pheromones for Integrated Pest Management in Vietnam. Integrated Pest Management (IPM): Environmentally Sound Pest Management 5: 103-127.
- Dey N, Biswas S, Ray Chaudhury T, Dey S R, De M, Ghose T K. 2005. RAPD based genetic diversity analysis of aromatic rice (*O. sativa* L.). Plant Cell Biotechnology and Molecular Biology 6: 133-142.
- Dey S R. 2016. Green Leafhopper (GLH), *Nephotettix virescens* (Distant) and Rice Tungro Disease (RTD). The Beats of Natural Sciences 3(3-4): 1-7.
- Dhaliwal G S, Jindal V, Dhawan A K. 2010. Insect pest problems and crop losses: Changing trends. Indian Journal of Ecology 37(1): 1-7.
- Dougoud J, Toepfer S, Bateman M, Jenner W H. 2019. Efficacy of homemade botanical insecticides based on traditional knowledge. A review. Agronomy for Sustainable Development. 39(37): 1-22.
- Dutta B C, Hazarika L K. 1994. Screening of some rice cultivars for resistance to rice hispa, *Dicladispa armigera* (Olivier) (Coleoptera: Chrysomelidae). Journal of Agricultural Science Societies 7(1): 31-34.
- Dutta S, Roy N. 2016. Life table and population dynamics of a major pest, *Leptocorisca acuta* (Thunb.) (Hemiptera: Alydidae), on rice and non-rice system. International Journal of Pure and Applied Bioscience 4(1): 199-207.
- Dutta S, Hossain A, Roy N. 2020. Population Ecology and Economic Thresholds for Sustainable Management of *Leptocorisca acuta* (Thunb.) on Three Rice Cultivars. Annals of Entomology 38(01-02): 55-76.
- Dutta S, Hossain A, Roy N. 2021. Rice cultivar selection against *Leptocorisca acuta* (thunb.) and their economic thresholds based ecological management. Journal of Entomological Research 45 (3): 405-414.
- Dutta S, Roy N. 2018a. Host preference and population dynamics of a major pest, *Leptocorisca acuta* (Thunb.) for their ecological management. International Journal of Biological Research 3: 31-39.
- Dutta S, Roy N. 2018b. Population dynamics and host preference of a major pest, *Scirpophaga incertulas* Walker (Pyralidae: Lepidoptera). International Journal of Advanced Science and Research 3: 120-127.
- Dutta S, and Roy N. 2022. Review on bionomics and management of rice stem borer. Journal of Entomology and Zoology Studies 10(5): 301-310.
- Gangwar R K. 2015. Life cycle and abundance of rice leaf folder, *Cnaphalocrocis medinalis* (Guenee)-A review. Journal of Natural Sciences Research 5 (15): 103-105.
- Gao X, Guo C, Li M, Li R, Wu X, Hu A, Hu X, Mo F, Wu S. 2020. Physicochemical properties and bioactivity of a new guar gum-based film incorporated with citral to brown planthopper, *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae). Molecules 25(9): 1-15.
- Ge L Q, Huang B, Jiang Y P, Gu H T, Xia T, Yang G Q, Liu F, Wu J C. 2017. Carboxylesterase precursor (EST-1) mediated the fungicide jinggangmycin-suppressed reproduction of *Sogatella furcifera* (Hemiptera: Delphacidae). Journal of Economic Entomology 110(5): 2199-2206.
- Ghosh A, Bandyopadhyay P, Mondal D, Shamurailatpam D, Karmakar A, Ghosh RK. 2015. Estimation of losses due to pests in rapeseed (*Brassica campestris*). In Weed Science for Sustainable Agriculture, Environment, and Biodiversity. (Eds. Rao A.N. and Yaduraju N.T.), Proceedings of 25th Asian Pacific Weed Science Society Conference, Hyderabad, Andhra Pradesh. India. 571 pp.
- Ghule S D, Patel K G, Pandya H V. 2008. Seasonal incidence of rice earhead bug (*Leptocorisca acuta*) of paddy in south Gujarat. Insect Environment 14(1): 7-8.
- Gunawardena N E, Bandumathie M K. 1993. Defensive secretion of rice bug, *Leptocorisca oratorius* Fabricius, (Hemiptera: Coreidae): A unique chemical combination and its toxic, repellent, and alarm properties. Journal of Chemical Ecology 19(4): 851-861.
- Gunawardena N E, Ranatunga R P. 1989. Laboratory and field studies of a natural attractant of the rice pest, *Leptocorisca acuta* (Hemiptera: Coreidae). Tropical Pest Management 35(2): 210-211.
- Gupta K, Kolhekar S, Paikra M, Nidhi. 2019. Cost benefit ratio for the management of rice gundhi bug, *Leptocorisca acuta* (Thonberg) under field condition. Journal of Pharmacognosy and Phytochemistry 2: 219-221.
- Hajjar M J, Ahmed N, Alhudaib K A, Ullah H. 2023. Integrated Insect Pest Management Techniques for Rice. Sustainability 15(5): 4499.
- Harborne J B. 1973. Methods of Plant Analysis. Phytochemical Methods 1: 1-32.
- Havelka J, Zemek R. 1999. Life table parameters and oviposition dynamics of various populations of the predacious gall-midge *Aphidoletes aphidimyza*. Entomologia Experimentalis et Applicata 91(3): 483-486.
- Huang X, Jiang T, Wu Z, Zhang W, Xiao H. 2020. Overwintering parasitism is positively associated with population density in diapausing larvae of *Chilo suppressalis*. Journal of Integrative Agriculture 19(3): 785-792.
- Huang Y, Jiang C, Liu C, Liu H, Yang Q, Wang H, Zeng W, Ouyang J, Xu K, Li Q. 2019. Overall evaluation of resistance capability of different rice varieties to *Sogatella furcifera* (Hemiptera: Delphacidae) using radar charts. Euphytica 215(58): 1-15.
- Hurali S, Biradar R, Vinoda H M, Masthanareddy B G, Gowdar S B. 2020. Bio-efficacy of new formulation of carbofuran against major insect pests of rice *Oryza sativa* L. International Journal of Chemical Studies 8(1): 1030-1035.
- Iqbal M F, Farooq M, Ahmad M, Latif M, Ahmad Z, Ali M A. 2015. Incidence of Rice Hispa (*Dicladispa armigera*) on paddy crop in Agro-Ecological Zone of District Sialkot. International Journal of Advanced Research in Biological Sciences 2(9): 64-66.
- January B, Rwegasira G M, Tefera, T. 2020. Rice stem borer species in Tanzania: a review. The Journal of Basic and Applied Zoology 81(1): 36.

- Jeevanandham N, Ramiah N, Chockalingam V, Jegadeesan R. 2020. An Overview of the Bionomics, Host Plant Resistance and Molecular Perspectives of *Sesamia inferens* Walker in Cereals and Millets. *Agronomy* 10(1705): 1-18.
- Jha S K., Monobrullah M, Singh P K, Prasad R. 2023. Performance of aromatic rice varieties against yellow stem borer *Scirpophaga incertulas*. *Indian Journal of Entomology* pp. 1-3.
- Jian X, Qin L, Chuan-ming L, Guang-jie H. 2019. Field effect of *Cnaphalocrocis medinalis* granulo virus (CnmeGV) on the pest of rice leafhopper. *Journal of Integrative Agriculture* 18(9): 2115-2122.
- Kakde A M, Patel K G. 2018. Screening of rice varieties against earhead bug, *Leptocorisa acuta* stal. (Hemiptera: Coreidae). *International Journal of Agriculture Sciences* 10: 5223-5226.
- Kakde A M, Patel K G, Tayade S. 2014. Role of life table in insect pest management-a review. *IOSR Journal of Agriculture and Veterinary Science* 7(1): 40-43.
- Katti G, Padmavathi C, Kumar R M. 2023. Integrated pest management in rice-based cropping systems in India. In *Integrated pest management in diverse cropping systems*. Apple Academic Press. pp. 111-135.
- Kaur R, Suri K S, Singh G. 2015. Influence of rice varieties on behaviour and reproductive physiology of *Sogatella furcifera* Horvath. *Agricultural Research Journal* 52(2):143-147.
- Keary M. 2023. A green theory of technological change: Ecologism and the case for technological scepticism. *Contemporary Political Theory*, 22(1): 70-93.
- Khalifa A A A A, El-Rewainy IM. (2012) Study some physiological characters, yield and yield component for five new rice varieties under different sowing dates. *Advances in Applied Science Research* 3(1): 440-445.
- Khare V K, Prakash R. 2020. Effect of abiotic factors on the population dynamics of paddy earhead bug, *Leptocorisa oratorius* F. *Journal of Entomology and Zoology Studies* 8(6): 157-160.
- Khatiwada J R, Ghimire S, Khatiwada S P, Paudel B, Bischof R, Jiang J, Haugaaseng T. 2016. Agriculture, ecosystems and environment frogs as potential biological control agents in the rice fields of Chitwan, Nepal. *Agriculture, Ecosystems and Environment* 230: 307-314.
- Kumar G S, Sarada O. 2020. Evaluation of Cow Based Fermented Organic Products for Non-insecticidal Pest Management in Castor. *International Journal of Current Microbiology and Applied Sciences* 9(10): 292-300.
- Kumar H. 2019. Resistance in rice against Brown planthopper, *Nilaparvata lugens* (Homoptera: Delphacidae) and White backed planthopper, *Sogatella furcifera* (Horvath) (Homoptera: Delphacidae): Role of new sources of resistance, BPH populations and screening methodology. *Journal Biological Series* 2(1): 001-026.
- Kumar J, Ramlal A, Mallick D, Mishra V. 2021. An overview of some biopesticides and their importance in plant protection for commercial acceptance. *Plants* 10(1185): 1-15.
- Kumar L V, Patil S U, Chakravarthy A K. 2012. Biochemical basis of resistance in rice against Asian rice gall midge, *Orseolia oryzae* (Wood-Mason) (Diptera: Cecidomyiidae). *Current Biotica* 6(2): 163-170.
- Kumari S, Yadav M, Prasad R, Prasad R, Kumar U. 2020. Effect of chemical insecticides on the incidence of gall midge (*Orseolia oryzae* Wood mason) in terms of silver shoots (SS%). *Journal of Entomology and Zoology Studies* 8(6): 781-785.
- Kunal G, Kumar R, Das R, Golda S. 2020. Life cycle of gundhi bug *Leptocorisa oratorius* F on alternate host *Echinochloa colonum* L under laboratory condition. *Journal of Pharmacognosy and Phytochemistry* 9(1): 543-546.
- Liu B, Syu K J, Zhang Y X, Gupta S, Shen Y J, Chien W J, Tseng J C, Lou D W. 2020. Practical Synthesis and Field Application of the Synthetic Sex Pheromone of Rice Stem Borer, *Chilo suppressalis* (Lepidoptera: Pyralidae). *Journal of Chemistry* 2020: 1-9.
- Longkumer I Y, Misra A K. 2020. Efficacy of essential oils against rice insect pest in Pusa, Bihar. *Journal of Entomology and Zoology Studies*. 8(3): 1980-1983.
- Luo D, Xia F, He M, Wu S, Zhao X, Liao X. 2023. Sublethal effects of the cis-nitromethylene neonicotinoid insecticide cyclopyrid on the white-backed planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae). *Crop Protection* 166(2023): 106172.
- Luo G H, Yao J, Yang Q, Zhang Z C, Hoffmann A A, Fang J C. 2016. Variability in development of the striped rice borer, *Chilo suppressalis* (Lepidoptera: Pyralidae), due to instar number and last instar duration. *Scientific Reports* 6(1): 1-7.
- Mandal K G, Thakur A K, Mohanty R K, Mishra A, Sinha S, Biswas B. 2022. Policy perspectives on agricultural water management and associated technologies suitable for different agro-climatic zones of West Bengal, India. *Current Science*. 122(4): 386.
- Manikandan N, Kennedy J S, Geethalakshmi V. 2015. Effect of temperature on life history parameters of brown planthopper (*Nilaparvata lugens* Stal). *African Journal of Agricultural Research* 10(38): 3678-3685.
- Matharu K S, Tanwar P S. 2020. Efficacy of different insecticides against brown planthopper, *Nilaparvata lugens* (Stål) in rice. *International Journal of Chemical Studies* 8(3): 870-873.
- Mathur K C, Rajamani S. 1984. *Orseolia* and rice: cecidogenus interactions. *Proceedings: Animal Sciences* 93(4): 283-292.
- Mobarak S H, Roy N, Barik A. 2020. Two-sex life table and feeding dynamics of *Spilosoma obliqua* Walker (Lepidoptera: Arctiidae) on three green gram cultivars. *Bulletin of Entomological Research* 110(2): 219-230.
- Minab F, Rakhshani E, Talamas E J, Moghaddam M G. 2023. A checklist of Platygasteridae and Scelionidae (Hymenoptera, Platygasteroidea) of Iran. *Journal of Insect Biodiversity and Systematics* 9(2): 343-383.
- Morya G P, Kumar R. 2019. Screening of rice germplasm resistance against major insect pests under eastern Uttar Pradesh conditions. *Journal of Entomology and Zoology Studies*. 7(6): 706-712.
- Nacro S, Heinrichs E A, Dakouo D. 1996. Estimation of rice yield losses due to the African rice gall midge, *Orseolia oryzivora* Harris and Gagne. *International Journal of Pest Management* 42(4): 331-334.
- Nathan S S, Kalaivani K, Murugan K, Chung P G. 2005. Efficacy of neem limonoids on *Cnaphalocrocis medinalis* (Guenée) (Lepidoptera: Pyralidae) the rice leafhopper. *Crop Protection* 24(8): 760-763.
- Ogah E O, Nwilene F E. 2017. Incidence of Insect Pests on Rice in Nigeria: A Review. *Journal of Entomology* 14: 58-72.
- Ouattara D, Nacro S, Latévi K, Coulibaly A, Somda I. 2020. Ecology of the African Rice Gall Midge, *Orseolia oryzivora* (Diptera: Cecidomyiidae) in Western Burkina Faso, West Africa. *Advances in Entomology* 8: 1-14.
- Padhan S, Raghuraman M. 2018. Bio-Efficacy of newer insecticides against Rice Gundhi Bug, *Leptocorisa acuta* (Thunberg) in Varanasi region. *Journal of Pharmacognosy and Phytochemistry* 7(3): 2068-2071.
- Padmavathi C, Katti G, Padmakumari A P, Voleti S R, Subba Rao L V. 2012. The effect of leafhopper *Cnaphalocrocis medinalis* (Guenee)

- (Lepidoptera: Pyralidae) injury on the plant physiology and yield loss in rice. *Journal of Applied Entomology* 137(4): 249-256.
- Padmavathi C, Katti G, Padmakumari P A, and Pasalu I C. 2008. Mortality and fertility life tables of leaf folder *Cnaphalocrosis medinalis* Guenee (Lepidoptera: Pyralidae) in rice. *Entomol* 33(1): 1-13.
- Pathak M D, Khan Z R. 1994. Insect pests of rice. International Rice Research Institute, Los Baños, Philippines. pp. 1-89.
- Pedigo L P, Higley L G. 1992. A new perspective of the economic injury level concept and environmental quality. *American Entomologist* 38: 12-21.
- Pedigo L P, Hutchins S H, Higley L G. 1986. Economic injury levels in theory and practice. *Annual Review of Entomology* 31: 341-368.
- Purnima D, Hazarika L K, Bora D, Puzari K C, Surajit K. 2013. Influence of storage conditions on viability and infectivity of talc based WP formulation of *Beauveria bassiana* against rice hispa, *Diadisa armigera* (Olivier). *Journal of Biological Control* 27(3): 229-233.
- Rahaman C R, Arko P S, Ali M E, Khan M A I, Apon S H, Nowrin F, Wasif A. 2020. Identification and recognition of rice diseases and pests using convolutional neural networks. *Biosystems Engineering* 194: 112-120.
- Rahaman M M, Islam K, Jahan M, Mamun M. 2014. Relative abundance of stem borer species and natural enemies in rice ecosystem at Madhupur, Tangail, Bangladesh. *Journal of the Bangladesh Agricultural University* 12(2): 267-272.
- Raj R S, Muthukrishnan N, Sathiah N, Prabakar K. 2020. Baseline susceptibility of different population of whitebacked planthopper, *Sogatella furcifera* from rice growing areas of Tamil Nadu, India to imidacloprid 17.8% S L. *Journal of Entomology and Zoology Studies* 8(5): 1894-1898.
- Rath P C, Bose L K, Subudhi H, N. Lenka, S. and Jambhulkar, N. N. (2020a) Biodiversity of insect pests of rice in India. *International Journal of Chemical Studies*. 8(1): 2998-3002.
- Rath P C, Bose L K, Subudhi H, Lenka S, Jambhulkar N N. 2020b. Biodiversity of Pests of Rice in Odisha. *International Journal of Current Microbiology and Applied Sciences*. 9 (3): 566-569.
- Raveau R, Fontaine J, Lounès-Hadj Sahraoui A. 2020. Essential oils as potential alternative biocontrol products against plant pathogens and weeds: A review. *Foods* 9(3): 1-31.
- Reji G, Chander S. 2008. A degree-day simulation model for the population dynamics of the rice bug, *Leptocoris acuta* (Thunb.). *Journal of Applied Entomology* 132(8): 646-653.
- Rothschild G H L. 1970. Observations on the ecology of the rice-ear bug *Leptocoris oratorius* (F.) (Hemiptera: Alydidae) in Sarawak (Malaysian Borneo). *Journal of Applied Ecology* 7(1): 147-394.
- Roy N. 2017. Life table and nutritional ecology of *Epilachna vigintioctopunctata* Fab. (Coleoptera: Coccinellidae) on three host plants. *International Journal of Horticulture* 7(2): 7-19.
- Roy N. 2019a. Jute leaf physicochemical cues mediated behavioural responses of *Diacrisia casignetum* Kollar. *Agricultural Research* 8: 287-296.
- Roy N. 2019b. Life table and economic threshold concept for ecologically sustainable management of *Diacrisia casignetum* Kollar (Lepidoptera: Arctiidae) on jute. *Entomol* 44(2): 103-110.
- Roy N. 2020. Population ecology and ETs based time series for climate smart pest management of *Spilosoma obliqua* Walker. *Entomol* 45(1): 15-30.
- Roy N. 2021. Population dynamics and economic thresholds based time series for smart pest management of sesame. *International Journal of Tropical Insect Science* 41: 2573-2584.
- Roy N. 2022. Population ecology and economic thresholds-based time series for climate smart pest management of *Spilosoma obliqua* Walker (Lepidoptera: Arctiidae) on three sesame cultivars. *The Journal of Basic and Applied Zoology* 83(1): 1-13.
- Sabale J P, Das C, Samui R P. 2010. Influence of weather factors on light trap catches of green leaf hopper at Pattambi, Kerala. *Journal of Agrometeorology* 12(1): 108-110.
- Sanchit S M, Sanchit S. 2019. Studies on host plant resistance against rice ear head bug, *Leptocoris acuta* Thunberg, Master Degree theses, Navsari Agricultural University, Navsari, Gujarat, India. 996 pp.
- Sankar S S H, Rani R P O. 2018. Pathogenicity and field efficacy of the entomopathogenic fungus, *Lecanicillium saksenae* Kushwaha, Kurihara and Sukarno in the management of rice bug, *Leptocoris acuta* Thunberg. *Journal of Biological Control* 32(4): 230-238.
- Sarao P S, Bentur J S. 2018. Quantification of antibiosis levels in nine different rice genotypes against *Nilaparvata lugens* (Homoptera: Delphacidae). *International Journal of Tropical Insect Science* 38(4), 330-339.
- Sarkar S K, Patra B C, Jana K, Ghosh K. 2016. Impact of different herbicides and their combinations on production economics of winter rice in West Bengal. *Indian Journal of Weed Science* 48(4): 442-444.
- Schoonhoven L M, Van Loon J J A, Dicke M. 2005. *Insect-plant biology*. Oxford University Press, Oxford, United Kingdom.
- Seleiman M F, Al-Suhaibani N, Ali N, Akmal M, Alotaibi M, Refay Y, Dindaroglu T, Abdul-Wajid, H H, Battaglia M L. 2021. Drought stress impacts on plants and different approaches to alleviate its adverse effects. *Plants* 10(2): 1-25.
- Selvaraj K, Chander S, Sujithra M. 2012. Determination of multiple-species economic injury levels for rice insect pests. *Crop Protection* 32: 150-160.
- Sharma R, Sharma P. 2021. Fungal entomopathogens: a systematic review. *Egyptian Journal of Biological Pest Control*, 31: 1-13.
- Shitiri A, Jamir I T, Neog P. 2014. Effect of cultivars and botanicals on the incidence of major insect pests in lowland rice. *International Journal of Bio-resource and Stress Management* 5(1): 58-63.
- Shivasharanappa Y, Pradeep S, Kumar M A A, Venkatesh H, Siddalingappa R. 2010. Biology of sugarcane internode borer, *Chilo sacchariphagus indicus* (Kapur). *Karnataka Journal of Agricultural Sciences*, 23(1): 140-141.
- Shyamrao I D, Raghuraman M. 2019. Bio-efficacy of insecticides against yellow stem borer (*Scirpophaga incertulas* Walker) in rice (*O. sativa* L.) ecosystem of Varanasi region. *Journal of Pharmacognosy and Phytochemistry* 8(2): 301-304.
- Song X Y, Peng Y X, Wang L X, Ye W N, Pei X G, Zhang Y C, Zhang S, Gao C F, Wu S F. 2022. Monitoring, cross-resistance, inheritance, and fitness costs of brown planthoppers, *Nilaparvata lugens*, resistance to pymetrozine in China. *Pest Management Science* 78(10): 3980-3987.
- Suzuki Y, Sogawa K, Seino Y. 1996. Ovicidal Reaction of Rice Plants against the Whitebacked Planthopper, *Sogatella furcifera* Horvath (Homoptera: Delphacidae). *Applied Entomology and Zoology* 31(1): 111-118.
- Tandon V, Srivastava A. 2019. Studies on life cycle of white stem borer, *Scirpophaga fusciflua* Hampson under Mid Hill conditions of Himachal Pradesh. *Oryza* 56(1): 33-37.
- Thein H W, Yamagata Y, Van Mai T, Yasui H. 2019. Four resistance alleles derived from *Oryza longistaminata* (A. Chev. & Roehrich) against green rice leafhopper, *Nephotettix cincticeps* (Uhler) identified using novel introgression lines. *Breeding Science* 69(4): 573-584.

- Tigga V, Kumar A, Sahu S P, Khan H H, Naz H. 2018. Assessment of the efficacy of certain chemical insecticides against rice gundhi bug, *Leptocorisca acuta* (Thun.) in Naini, Allahabad region. International Journal of Chemical Studies 6(1): 959-961.
- Win S S, Muhamad R, Ahmad Z, Adam N. 2011. Life Table and Population Parameters of *Nilaparvata lugens* Stal. (Homoptera: Delphacidae) on Rice. Tropical Life Sciences Research 22: 25-35.
- Win S S, Muhamad R, Ahmad Z A M, Adam N A. 2009. Life table and population parameter of *Sogatella furcifera* (Horvath) (Homoptera: Delphacidae) on rice. Journal of Biological Sciences 9: 904-908.
- Xu J, Wang X, Han C, Jiang Y, Zhong W, Liu B. 2022. Trait-based comparison of transgenic Bt rice and its non-Bt counterpart in response to soil copper pollution. Environmental Science and Pollution Research, 29: 2773-2782.
- Yuan P L, Wang J P, Can G U O, Guo Z Y, Yao G U O, Cao C G. 2022. Sustainability of the rice-crayfish farming model in waterlogged land: A case study in Qianjiang County, Hubei Province, China. Journal of Integrative Agriculture 21(4): 1203-1214.
- Zhao X, Li C, Xu J, Zhang X, Xiong H, Huang Q, Cao S. 2019. Field-scale evaluation on the efficacy of a novel benzoylphenylurea TXH09 against chewing insect pests harming vegetables and rice. Crop Protection 126: 1-6.

(Manuscript Received: December, 2022; Revised: May, 2023;

Accepted: May, 2023; Online Published: May, 2023)

Online First in www.entosocindia.org and indianentomology.org Ref. No. e23949