



HOST PREFERENCE AND POPULATION DYNAMICS OF *HOLOTRICHIA NAGPURENSIS* KHAN AND GHAI

MAYANK KUMAR* AND A K PANDEY

Department of Entomology, College of Agriculture,
G B Pant University of Agriculture and Technology, Pantnagar 263145, Uttarakhand, India

*Email: mayankkumar1411@gmail.com (corresponding author)

ABSTRACT

Holotrichia nagpurensis Khan and Ghai is a major white grub species of subfamily Melolonthinae. Its wide host range has been reported from different parts of India. An experiment was carried out to find host preference and population dynamics at three locations of Pantnagar in Terai region of Udham Singh Nagar District during 2018-19. Beetles were recorded from six host plants among which neem *Azadirachta indica*, was the major one. The p-value of two-way ANOVA between populations of three locations ($0.0006 < 0.001$) and from six hosts ($0.0002 < 0.001$) showed that there exists significant difference in distribution and feeding preference of *H. nagpurensis* on host plants. Among the hosts, *A. indica* was found to be the most preferred with maximum adult density (419 adults) and average 46.55 beetles/ tree; and the multiple comparisons revealed a significant host preference. Correlation coefficients revealed that minimum relative humidity exhibits a negative relationship with beetle emergence.

Key words: Scarabaeidae, *Holotrichia nagpurensis*, adults, host preference, *Azadirachta indica*, population dynamics, weather parameters, relative humidity, correlation coefficients

White grub, also known as May-June beetles, belonging to family Scarabaeidae is a major insect pest. Its adults are nocturnal, and feed on the leaves and soft shoot and fruits of various trees, shrubs and grasses (Ritcher, 1958; Vallejo et al., 1998). Their polyphagous nature make them major pests in India (Metcalf and Luckman, 1975). Of the 2000 species known from the Indian subcontinent 40 species cause serious damage to various crops (Veeresh et al., 1991). Among these, *Holotrichia* spp. (subfamily Melolonthinae) are mostly leaf feeders in adult stage (Arrow, 1917); this genus has >100 species with wide distribution (Mathur et al., 2010). There are species like *H. consanguinea*, *H. longipennis*, *H. serrata*, *H. insularis* etc observed from 27 host plants in north India (Srivastava and Khan, 1963; Bhadauria and Nigam, 1982; Haq, 1962). Many abiotic factors influence their distribution and diversity. High diversity of phytophagous insects is also the result of factors that affecting their diet breadth (Gaete-Eastman et al., 2004). There is no information about the host range and feeding preference of *H. nagpurensis* on host plants from Terai area of Pantnagar of Udham Singh Nagar district of Kumaon. This study evaluates the host range, feeding preference and population dynamics of its adults.

MATERIALS AND METHODS

The present study was conducted at three locations

i.e. Crop Research Centre (CRC), Horticulture research center (HRC) and Livestock Research Center (LRC) of Udham Singh Nagar in Kumaon region Uttarakhand during 2018-19. Weekly surveys were made to record the *H. nagpurensis* incidence on various host plants like neem, guava, jackfruit, litchi, mango, bakane, amaltash, teak, pride of India and ashok in addition to some cultivated field crops like maize, soybean, sugarcane, rice and calotropis. Weekly observations were made starting from appearance of beetle i.e. from 10th standard meteorological weeks (SMW) to 25th MSW (from March to June), with counting the of the adults feeding on leaves using powerful torch during night. The beetles were also collected from each host plants available on experimental site by shaking the branches to dislodge the beetles. Collected beetles were brought to the laboratory where, they were killed and sorted out before storing. Number of beetles that flew away from tree were also included. Because of largeness in size these were easily identified during flight (Litsinger et al., 2002). The data on the cumulative number of beetles for each tree species was calculated to evaluate the host preference. Data on weather parameters viz., weekly maximum and minimum temperature ($^{\circ}\text{C}$), relative humidity (RH) (%) at 7:12 am and 2:12 pm, rainfall (mm), wind velocity (km/ hr.) and sunshine (hr) were obtained. The data were subjected to statistical analysis by ANOVA and LSD test (Litsinger et al., 2002) using R

and SPSS software packages, respectively. Correlation coefficients of the incidence of beetle with weather factors were computed with R software.

RESULTS AND DISCUSSION

The results of the study brought out the host range/preference and population dynamics of *H. nagpurensis*. These included the major host plants like *A. indica*, *P. guajava*, *A. heterophyllum*, *L. scinensis*, *M. indica*, *M. azadirach*, *C. fistula*, *T. grandis*, *L. speciosa* and *Polyalthia* sp. in addition to some cultivated field crops like *Z. mays*, *G. max*, *S. officinarum*, *O. sativa* and *Calotropis* sp. These revealed the wide host range with significant variations and choice of host plant for feeding; significantly maximum (713 beetles) was observed at the location HRC followed by LRC (698) and CRC (503) on the preferred hosts. Among the 15 host plants selected, which are common to all the sites, six trees i.e. *A. indica*, *M. indica*, *A. heterophyllum*, *P. guajava*, *Z. mays* and *M. azadirach* inhabited maximum adults (Table 1); of these *A. indica* was the most preferred inhabiting >one fifth, and the least (11.91%) on *M. azadirach* followed by *Z. mays* (11.54%) only. On an average, *A. indica* recorded 46.55 adults/ tree. Two-way ANOVA revealed significant differences among the locations studied, and among the six host trees. The multiple comparison values also indicate that *A. indica* was significantly most preferred (Table 2). The emergence of *H. nagpurensis* started from 7 pm of 10th and up to 25th MSW with a peak during 16th MSW (Fig. 1); correlation coefficients revealed a non-significant but positive correlation with maximum temperature ($r=0.167$), and a negative one with minimum temperature ($r=-0.130$); negative relationship with both maximum and minimum RH and

Table 2. Multiple comparisons of hosts for feeding preference (2018)

S. No.	I-Sample	J- Sample	Mean Diff.	Sig.
1.	<i>A. indica</i> L.	<i>Z. mays</i> L.	66.00	.003*
		<i>P. guajava</i> L.	19.00	.316ns
		<i>A. heterophyllum</i> Lam.	20.00	.292ns
		<i>M. azadirach</i> L.	63.67	.004*
		<i>M. indica</i> L.	31.33	.110ns
2.	<i>Z. mays</i> L.	<i>A. indica</i> L.	-66.00	.003*
		<i>P. guajava</i> L.	-47.00	.024*
		<i>A. heterophyllum</i> Lam.	-46.00	.026*
		<i>M. azedarach</i> L.	-2.33	.900ns
		<i>M. indica</i> L.	-34.67	.080ns
3.	<i>P. guajava</i> L.	<i>A. indica</i> L.	-19.00	.316ns
		<i>Z. mays</i> L.	47.00	.024*
		<i>A. heterophyllum</i> Lam.	1.00	.957ns
		<i>M. azadirach</i> L.	44.67	.030*
		<i>M. indica</i> L.	12.33	.510ns
4.	<i>A. heterophyllum</i> Lam.	<i>A. indica</i> L.	-20.00	.292ns
		<i>Z. mays</i> L.	46.00	.026*
		<i>P. guajava</i> L.	-1.00	.957ns
		<i>M. azadirach</i> L.	43.67	.033*
		<i>M. indica</i> L.	11.33	.544ns
5.	<i>M. azadirach</i> L.	<i>A. indica</i> L.	-63.67	.004*
		<i>Z. mays</i> L.	2.33	.900ns
		<i>P. guajava</i> L.	-44.67	.030*
		<i>A. heterophyllum</i> Lam.	-43.67	.033*
		<i>M. indica</i> L.	-32.33	.100ns
6.	<i>M. indica</i> L.	<i>A. indica</i> L.	-31.33	.110ns
		<i>Z. mays</i> L.	34.67	.080ns
		<i>P. guajava</i> L.	-12.33	.510ns
		<i>A. heterophyllum</i> Lam.	-11.33	.544ns
		<i>M. azadirach</i> L.	32.33	.100ns

ns=non- significant; significant at $p=0.05$

Table 1. Incidence of *H. nagpurensis* on hosts in three locations (2018-19)

S. No.	Host name	CRC	HRC	LRC	Total	% of total
1	<i>Azadirachta indica</i> L.	115	147	157	419	21.89
2	<i>Zea mays</i> L.	55	98	68	221	11.54
3	<i>Psidium guajava</i> L.	85	139	138	362	18.91
4	<i>Atrocarpus heterophyllum</i> Lam.	89	142	128	359	18.75
5	<i>Melia azedarach</i> L.	64	77	87	228	11.91
6	<i>Mangifera indica</i> L.	95	110	120	325	16.98
	Total	503	713	698	1914	




Fig. 1. Seasonal emergence of *H. nagpurensis* vs. weather factors

rainfall were also observed of which only the one with minimum RH ($r=-550^*$) was significant. These results corroborate with those of Pal (1977) and Gupta (1973); and weather factors and availability of desirable host are important (Veeresh, 1988; Ratnadass et al., 2012). Present results are in partial agreement with those of Mishra and Singh (1999) on the favourable weather. Prathibha et al. (2013) also reported that rainfall is an important factor relating to emergence and aggregation of this beetle, and significant correlation with maximum temperature corroborates with that of Seram and Saikia (2015).

ACKNOWLEDGEMENTS

Authors thank Dr Arjun Singh Baloda, Coordinator, All India Network Project of White grub and Other Soil Arthropods for providing financial support; and in-charges of Crop Research Centre, Horticulture Research Centre and Livestock Research Centre, GBPUAT, Pantnagar for support.

REFERENCES

- Arrow G J. 1917. The fauna of British India. Including Ceylon and Burma. Coleoptera, Lamellicornia Part 2 (Rutelinae: Desmonycinae and Euchirinae). Taylor and Francis, London. 5(13): 1-187.
- Bhadauria A S, Nigam P M. 1982. Food preference to *Holotrichia consanguinea* (Coleoptera: Scarabaeidae). Entomon 7(3): 331-332.
- Gaete-Eastman C, Figueroa C C, Olivares-Donoso R, Niemeyer H M, Ramírez C C. 2004. Diet breadth and its relationship with genetic diversity and differentiation: the case of southern beech aphids (Hemiptera: Aphididae). Bulletin of Entomological Research 94(3): 219-227.
- Gupta K M. 1973. Neem leaves attract white grub beetles. Indian Journal of Entomology 35: (3) 276.
- Haq A. 1962. Notes on the Bionomics of *Lachnosterna longipennis* B1. (Melolonthinae: Coleoptera). Indian Journal of Entomology 24(3): 220-221.
- Litsinger J A, Libetario E M, Barrion A T. 2002. Population dynamics of white grubs in the upland rice and maize environment of Northern Mindanao, Philippines. International Journal of Pest Management 48(3): 239-260.
- Metcalf R L, Luckman W H. 1975. Introduction to pest management. John Wiley and sons, New York, NY. 587 pp.
- Mathur Y S, Bhatnagar A, Singh S. 2010. Bioecology and management of phytophagous white grubs of India. Technical Bulletin 4. All India Network Project on White grubs and Other Soil Arthropods, Agriculture Research Station, Durgapura, Jaipur.
- Mishra P N, Singh M P. 1999. Determination of predominate species of white grubs species in Garhwal region of Uttara Pradesh Hills (India). Journal of Entomological Research 23(1): 65-69.
- Pal S K. 1977. White grubs and their management, Monograph No. 5. Central Arid Zone Research Institute, Jodhpur.
- Prathiba P S, Kumar A R V, Subaharan K. 2013. Ethology of coconut root grub chafer *Leucopholis coneophora* Burmeister (Melolonthinae: Scarabaeidae). International Journal of Agriculture and Food Science Technology 4(2): 24-28.
- Ratnadass A, Fernandes P, Avelino J, Habib R. 2012. Plant species diversity for sustainable management of crop pests and diseases in agroecosystems: a review. Agronomy for Sustainable Development 32(1): 273-303.
- Ritcher P O. 1958. Biology of Scarabaeidae. Annual Review of Entomology 3(1): 311-334.
- Seram D, Saikia K, 2015. Weather correlation of white grub, *Leucopholis coneophora* (Burmeister) incidence in mid-hills of Meghalaya. Life Science International Research Journal 2(2): 286-288.
- Srivastava B K, Khan R M. 1963. Bionomics and control of *Holotrichia insularis* (Melolonthidae: coleoptera). Indian Journal of Entomology 25(4): 347-354.
- Vallejo F M A, Moron, Orduz S. 1998. First report and description of immature stages of *Phyllophaga obsoleta* (Blanchard) (Col: Melolonthidae) in Colombia. The Coleopterist's Bulletin 52(2): 109-117.
- Veeresh G K, 1988. White grubs. Applied soil biology and ecology. G K Veeresh, D Rajgopal (eds.) (2nd Ed) IBH, New Delhi. 243-282 pp.
- Veeresh G K, Kumar A R V, Ali M T M. 1991. Biogeography of pest species of white grubs of Karnataka. Veeresh G K, Rajagopal D, Viraktamath C A (eds.). Advances in management and conservation of soil fauna. Oxford and IBP Publishing Company Pvt. Ltd., Bangalore. 191-198 pp.

(Manuscript Received: November, 2020; Revised: January, 2021;

Accepted: January, 2021; Online Published: August, 2021)

Online published (Preview) in www.entosocindia.org Ref. No. e20398